

SPECIAL ARTICLES

Consensus Statement of the American Academy of Sleep Medicine on the Recommended Amount of Sleep for Healthy Children: Methodology and Discussion

Shalini Paruthi, MD, Moderator¹; Lee J. Brooks, MD^{2,3}; Carolyn D'Ambrosio, MD⁴; Wendy A. Hall, PhD, RN⁵; Suresh Kotagal, MD⁶; Robin M. Lloyd, MD⁶; Beth A. Malow, MD, MS⁷; Kiran Maski, MD⁸; Cynthia Nichols, PhD⁹; Stuart F. Quan, MD¹⁰; Carol L. Rosen, MD¹¹; Matthew M. Troester, DO¹²; Merrill S. Wise, MD¹³

¹Saint Louis University, St. Louis, MO; ²Children's Hospital of Philadelphia, Philadelphia, PA; ³Liaison for the American Academy of Pediatrics; ⁴Brigham & Women's Hospital, Boston, MA; ⁵University of British Columbia School of Nursing, Vancouver, BC; ⁶Mayo Clinic, Rochester, MN; ⁷Vanderbilt University Medical Center, Nashville, TN; ⁸Boston Children's Hospital, Boston, MA; ⁹Munson Sleep Disorders Center, Traverse City, MI; ¹⁰Harvard Medical School, Boston, MA; ¹¹Rainbow Babies & Children's Hospital, Cleveland, OH; ¹²Barrow Neurologic Institute at Phoenix Children's Hospital, Phoenix, AZ; ¹³Methodist Healthcare Sleep Disorders Center, Memphis, TN

Members of the American Academy of Sleep Medicine developed consensus recommendations for the amount of sleep needed to promote optimal health in children and adolescents using a modified RAND Appropriateness Method. After review of 864 published articles, the following sleep durations are recommended: Infants 4 months to 12 months should sleep 12 to 16 hours per 24 hours (including naps) on a regular basis to promote optimal health. Children 1 to 2 years of age should sleep 11 to 14 hours per 24 hours (including naps) on a regular basis to promote optimal health. Children 3 to 5 years of age should sleep 10 to 13 hours per 24 hours (including naps) on a regular basis to promote optimal health. Children 6 to 12 years of age should sleep 9 to 12 hours per 24 hours on a regular basis to promote optimal health. Teenagers 13 to 18 years of age should sleep 8 to 10 hours per 24 hours on a regular basis to promote optimal health. Sleeping the number of recommended hours on a regular basis is associated with better health outcomes including: improved attention, behavior, learning, memory, emotional regulation, quality of life, and mental and physical health. Regularly sleeping fewer than the number of recommended hours is associated with attention, behavior, and learning problems. Insufficient sleep also increases the risk of accidents, injuries, hypertension, obesity, diabetes, and depression. Insufficient sleep in teenagers is associated with increased risk of self-harm, suicidal thoughts, and suicide attempts.

Commentary: A commentary on this article appears in this issue on page 1439.

Keywords: pediatric, sleep duration, consensus

Citation: Paruthi S, Brooks LJ, D'Ambrosio C, Hall WA, Kotagal S, Lloyd RM, Malow BA, Maski K, Nichols C, Quan SF, Rosen CL, Troester MM, Wise MS. Consensus statement of the American Academy of Sleep Medicine on the recommended amount of sleep for healthy children: methodology and discussion. *J Clin Sleep Med* 2016;12(11):1549–1561.

1.0 INTRODUCTION

Healthy sleep requires adequate duration, appropriate timing, good quality, regularity, and the absence of disturbances and disorders. Sleep duration is a frequently investigated sleep measure in relation to health outcomes. Many studies have shown that adequate sleep duration is associated with better attention, behavior, cognitive functioning, emotional regulation, and physical health among children.^{1–3}

A panel of 13 experts in sleep medicine and research used a modified RAND Appropriateness Method⁴ to develop recommendations regarding the sleep duration range that promotes optimal health in children aged 0–18 years. The Consensus Recommendations⁵ were previously published, and this report expands on the methodology, results of the literature search, key findings from published research, overview of discussion by the panel members during the in-person meeting, and limitations of the process to answer the critical question: How much sleep is needed for optimal health in children?

2.0 METHODS

The American Academy of Sleep Medicine (AASM) Sleep Duration Consensus Conference used a modified RAND Appropriateness Method (RAM)⁴ to establish consensus for the amount of sleep needed to promote optimal health in children and teenagers.

2.1 Expert Panel Selection

In accordance with the recommendations of the RAM, the Sleep Duration Consensus Conference panel comprised 13 voting members, including a moderator. All panel members are experts in sleep medicine and/or sleep science. The panel members were recommended by the Board of Directors of the AASM.

Panel members were sent a formal letter of invitation from the AASM and were required to complete Conflict of Interest disclosures before being officially accepted. To avoid further conflicts, panel members were not permitted to participate in similar consensus activities by other organizations.

Table 1—Data extracted from studies for evidence tables.

1	Study design
2	Oxford grade
3	Number of study participants
4	% of female participants
5	Method by which sleep duration was obtained
6	Age range of child participants
7	Major outcomes and conclusions

2.2 Modified RAND Appropriateness Method

The RAND Appropriateness Method uses a detailed search of the relevant scientific literature, followed by two rounds of anonymous voting to determine consensus on the appropriateness of a recommendation. The first round of voting is completed without panel interaction to prevent panel members from influencing each other's votes. The second round of voting occurs after a panel discussion of the available evidence and Round 1 voting results.

In a modification to RAM, the Consensus Conference included a third round of voting, which considered all available evidence and the previous voting results, to establish a single recommendation for the amount of sleep needed to promote optimal health for each age group in children. The third round also involved a discussion of the merits of recommending an optimal sleep duration range versus a simple threshold value. The final Consensus Recommendations resulted from the third round of voting.

The charge to the Consensus Conference panel was to determine a sleep duration recommendation for healthy children. Panel members voted on the appropriateness of one-hour increments ranging from < 6 to ≥ 18 hours of sleep. One-hour increments were selected because these were the most commonly reported units in epidemiologic and experimental studies. Substantial heterogeneity was present in the sleep duration assessment instruments. The consensus recommendations focused on overnight and daytime nap durations when appropriate as napping is considered biologically normal under the age of 7 years. The final recommendations were based on the one-hour values that were determined by the panel to be "appropriate" to promote optimal health in children.

2.3 Detailed Literature Search and Review

The AASM Board of Directors charged the panel with developing a recommendation for sleep duration in healthy children. This charge coincides with the goals of the National Healthy Sleep Awareness Project (NHSAP) and with the Sleep Health Objective of Healthy People 2020 to "increase the proportion of students in grades 9 to 12 who get sufficient sleep."⁶

The scope of the recommendation was limited to children aged 0–18 years. After a preliminary review of the literature, prior Centers for Disease Control and Prevention (CDC), AASM, and National Sleep Foundation (NSF) recommendations, as well as commonly frequented websites, the following age groups were created: < 12 months, 12 months to < 3 years, 3 years to < 6 years, 6 years to < 13 years, and 13–18 years. There was substantial overlap of age groups within the literature, and this was discussed during the in-person conference.

Table 2—Health domains.

1	Cardiovascular health
2	Developmental health
3	Human performance
4	General health
5	Immunology
6	Longevity
7	Mental health
8	Metabolic health
9	Cancer
10	Pain

A preliminary search of the literature and specific National Library of Medicine Medical Subject Headings (MeSH) terms identified several health outcomes that were most commonly examined in relation to sleep duration in children. Based on this evidence, the panel decided to focus on the relationships between sleep duration and the following 10 health categories: cardiovascular health, developmental health, human performance, general health, immunology, longevity, mental health, metabolic health, pain, and cancer.


After establishing the health categories, a detailed literature search was performed in PubMed on October 1, 2015. The search terms used for the literature search are detailed in the supplemental material. The search was restricted to studies in human children ages 0–18 years, published in English, with no publication date limit. Case reports, editorials, commentaries, letters and news articles were excluded from the search results. The initial search produced 1,040 publications. The search results were reviewed based on title and excluded *a priori* for the following reasons: focus on sleep quality or fatigue instead of sleep duration, assessing sleep duration in specific disorders or sleep disorders, experiments on total sleep deprivation, children sleeping outside normal day/night sleep schedules, assessments of sleep deprivation as a treatment (insomnia or depression), and focusing on medication effects on sleep duration. Application of these restrictions resulted in 864 publications.

The panel reviewed the abstracts of these remaining publications using the criteria described above. Citation pearl growing was used to capture additional important publications that were not identified by the search. Accepted publications were graded for quality using the Oxford criteria.⁷ All accepted publications were reviewed in detail and the data listed in **Table 1** were extracted.

Based on the data extraction, accepted studies were subdivided into the categories listed in **Table 2**.

The extraction sheet and full text of all accepted publications were made available to the panel members for review. Each panel member was assigned to a particular health domain and asked to identify the most informative studies based on study design and evidence quality. After review of the literature, the domains of pain and cancer did not contain sufficient data to guide sleep duration recommendations and therefore were excluded prior to Round 1 Voting. After lengthy discussion at the conference, the domain of longevity was excluded prior to Round 2 Voting. A second PubMed literature search

Figure 1—Round 1 voting results.



Round 1 voting was based on voting across all age groups under the following eight domains: cardiovascular health, developmental health, human performance, general health, immunology, longevity, mental health and metabolic health.

was performed during the conference on February 19, 2016 to collect more recent relevant studies.

2.4 Round 1 Voting

Prior to the conference, panel members reviewed the accepted publications and extraction sheets. Based on their review of this material and their clinical and research expertise, members voted to indicate their agreement with the following statement, “Based on the available evidence, [x] hours of sleep is associated with optimal health within the [x] category in the [x] age group.” “Hours of sleep” was categorized as: < 6 hours, 6 to < 7 hours, 7 to < 8 hours, 8 to < 9 hours, 9 to < 10 hours, 10 to < 11 hours, 11 to < 12 hours, 12 to < 13 hours, 13 to < 14 hours, 14 to < 15 hours, 15 to < 16 hours, 16 to < 17 hours, 17 to < 18 hours, ≥ 18 hours. The panel members voted using a 9-point Likert scale, where 1 meant “strongly disagree,” 9 meant “strongly agree,” and 5 meant “neither disagree nor agree.” Panel median values were placed into three broader categories, with the following interpretations: 1–3 indicated disagreement with the statement, 4–6 indicated uncertainty, and 7–9 indicated agreement with the statement.

Panel members were instructed not to discuss the evidence or their votes with each other to ensure independence. Panel members’ votes were collected by email and compiled to

determine the median and distribution of votes. Individual results tables were created and distributed to the members at the consensus conference, displaying the distribution of votes (anonymously), the member’s vote, and the median vote (Figure 1).

2.5 Conference Proceedings and Round 2 Voting

Prior to the conference, panel members were selected to become domain experts for each domain. At the conference, members reviewed the results of Round 1 voting for a domain, and then the domain experts presented a review of the best available evidence for that domain. Panel members then discussed the results of Round 1 voting, the accepted publications for the domain and any other relevant evidence. After discussion, panel members completed Round 2 voting for the age groups within the domain, following the same procedures from Round 1 voting. The conference proceeded in this manner for each domain.

2.6 Round 3 Voting and Development of Recommendations

Panel members reviewed and discussed Round 2 voting results for all domains and the entire body of accepted publications in preparation of voting on recommendation statements. After

discussions concluded, panel members completed Round 3 voting for a single recommendation of appropriate sleep duration range for each age group, following the same procedure as for Round 1 and Round 2 voting, but with the following statement: “Based on the available evidence, [x] hours of sleep is associated with optimal health in [x] age group.”

Based on the results of Round 1 and Round 2 voting, the conference discussions, and with the agreement of all the panel members, the infant (0 to 12 months) category was reduced to 4 to 12 months. This decision was based on the lack of evidence in this 0–4 month old age group. Thus, no recommendations were made for children under 4 months of age for any of the categories.

Upon completion of Round 3 voting, the panel members reviewed the voting results and developed the recommendations. The age groups and hour ranges were simplified to those presented in the consensus statements for the purposes of simplifying recommendations and ensuring clarity.

The final recommendations were submitted to the AASM, Sleep Research Society (SRS), American Academy of Pediatrics (AAP), and American Association of Sleep Technologists (AASST) Boards of Directors for their endorsements.

3.0 SUMMARY OF LITERATURE

The following sections summarize the key evidence considered by the panel in developing the recommendations while acknowledging that a complete evaluation of the evidence is beyond the scope of this consensus process.

For each domain, the panel reviewed studies with children from all over the world, addressing sleep duration and health outcomes across the prespecified age ranges. Within each domain, there were topics without information for some of the age groups, and often studies spanned several of our prespecified age groups. Studies that assessed the relationship between sleep duration and the search term of interest were the most informative. Many studies reported more than one outcome. However, emphasis was placed on longitudinal or cross-sectional cohort studies that reported sleep duration in unselected general populations as well as smaller studies of unconstrained sleep in healthy children. Some studies only provided correlation or regression coefficients and thus were minimally informative. Others reported the association between general health and sleep duration, but only at a limited number of thresholds. In most studies, sleep duration was assessed by parent or child self-report. Polysomnographic data and actigraphy were considered when appropriate such as in studies of unconstrained sleep in the laboratory or at home. Systematic comparisons between studies were challenging and in most cases not possible. The following domains and information were reviewed.

3.1 General Health

In the general health domain, the number of children in each study ranged from less than 25 to over 74,000.

Within the 4-months to < 12 months age range, there were few studies. However, two large prospective birth cohorts indicated that over a 24-hour period, healthy infants slept

slightly > 13 to slightly > 14 hours.^{8,9} Furthermore, infants sleeping for lesser amounts of time had a greater likelihood of quality of life issues later in childhood.⁹

For children in the 1 to < 3 years and 3 to < 6 years age ranges, there were only a small number of studies to review. One cohort study reported sleep durations between 11 and 12 hours in normal children.⁸ Limited data also suggested that sleeping less than 10 hours was associated with a greater risk of accidental injury¹⁰ and reduced quality of life several years later.¹¹

More studies were available that addressed general health outcomes in older children including those with polysomnographic data during unconstrained sleep.^{12,13} In the 6 to < 13 years age group, the panel observed that this was a period of rapid physical and mental development with more sleep appropriate for children at the lower end and less sleep for those at the higher end of the age range. Most informative was a large meta-analysis of children from 20 countries that indicated children between 9 and 12 years of age slept approximately 10 hours per night.¹⁴ In addition, data were available suggesting children sleeping 10 hours or more per night reported better health.¹⁵ In the 13–18 year age range, increased rates of injury were noted for those sleeping less than 7 or 8 hours^{16,17} and better health was reported for those sleeping 9 hours or more.¹⁵ A meta-analysis found that in teenagers, sleep declined continuously from between 9 and 10 hours at age 13 to between slightly less than 8 to slightly greater than 9 hours at age 18 years of age.¹⁵ Extended sleep duration was noted on weekends compared with weekdays.

In summary, limited information indicates that there is a continuous decline in the amount of sleep required for normal general health. Lesser amounts are associated with poorer overall health and reduced quality of life later in childhood. However, there is little information regarding any impact of excessive sleep on overall health.

3.2 Cardiovascular Health

Many of the studies in the cardiovascular domain examined the relationship between sleep duration and hypertension. Others looked at body mass index, waist circumference, serum lipids, C-reactive protein (CRP), and hemoglobin A1C (HbA1C). Most studies were cross-sectional, observational, retrospective or cohort design; none were randomized controlled studies. The number of participants varied from as few as 20 to just over 5,500. No studies in this domain included children younger than 5 years of age. Most studies focused on adolescents and teenagers.

The majority of studies suggested a shorter sleep duration was associated with either an increased risk of hypertension or actual hypertension. However, the definition of “short/shorter” sleep was different between the studies. For children > 5 years, studies suggested a shorter sleep duration (< 10 hours,¹⁸ some < 9 hours,¹⁹ some < 8 hours,^{20,21} and one < 5 hours²²) was associated with either an increased risk of hypertension or actual hypertension. One study showed an increased odds ratio of hypertension in girls who slept > 9.5 hours.²³ Another study suggested that < 10 hours of sleep was associated with hypotension (systolic blood pressure < 100 mm Hg) in preschoolers.¹⁸

Conversely, there were also data to suggest no adverse effect of sleep characteristics on cardiovascular health.^{24–26}

In summary, evidence indicates that shorter sleep duration of < 5 hours is associated with increased risk of hypertension. However, there is some suggestion that in girls regularly sleeping more than 9.5 hours, hypertension risk is increased.

3.3 Metabolic Health

Many studies in the metabolic domain specifically targeted the relationship between sleep duration and overweight/obesity. Other metabolic parameters studied included: waist circumference, insulin resistance (HOMA-IR), bone mineral content, triglycerides, C-reactive protein, HbA1c, leptin, and ghrelin. Most were cross-sectional studies of community-based populations. The number of participants in studies ranged from 62 to 81,390.

Most studies demonstrated a negative association between sleep duration and overweight/obesity. Shorter sleep durations were also associated with increased risk for developing overweight/obesity. Two meta-analyses found sleep duration inversely correlates with obesity in children 0–18 years.^{27,28} For each hour increase in sleep, the risk of overweight/obesity decreased. Children under age 10 years show a more linear dose-response relationship of sleep duration and weight status. Additionally, there is evidence of a stronger inverse relationship for sleep duration and weight status in boys compared with girls.²⁸

A longitudinal study of 915 children aged 6 months to 3 years, showed infant sleep duration of less than 12 hours per day was associated with higher body mass index (BMI) z scores and increased odds of overweight during subsequent preschool years.²⁹ Similarly, in a cohort study of 1,930 children ages 0–13 years, in younger children, ages 0–4 years, sleep duration of less than 11 hours was subsequently associated with increased risk of being overweight/obese. However for children ages 5–13 years, sleep duration was not associated with subsequent weight status.³⁰ Other cohort studies report similar findings.³¹ A large cross-sectional study of 3,875 infants and 3,844 children (up to age 7 years) showed that sleep duration did not predict obesity; however, obese children ages 6 to 7 years were found to sleep approximately 30 minutes less than non-obese children.³² In a cross-sectional study of 1,229 children ages 5 to 11 years, children who slept less than 10 hours per weeknight were 5 times more likely to be overweight than those who slept at least 12 hours on weeknights.³³ In another large cross-sectional study of 8,274 children ages 6 to 7 years old, children with sleep duration shorter than 10 hours had an increased odds ratio to develop obesity.³⁴ In a cross-sectional study of 6,576 children, ages 7–11 years, children who slept < 9 hours per night had a higher risk for overweight, obesity, and abdominal obesity compared to children who slept 10–11 hours.³⁵ In a study of 319 children ages 10–17 years, total sleep time was negatively correlated with overweight/obese status.³⁶ Similar findings were observed in a larger cross-sectional study of 6,324 children ages 7–15 years. Boys who slept < 10 hours per night had increased odds of overweight compared to children sleeping > 10 hours.³⁷ In 3,311 children 12.5 to 17.5 years, shorter sleep duration < 8 hours was associated with increased

BMI, body fat, and waist and hip circumferences.³⁸ A cohort study of 1,303 children aged 5–29 months showed that children sleeping < 10 hours per night consumed approximately 50 more kcal per day than children sleeping 11–12 hours,³⁹ suggesting that changes in appetite may be a possible mechanism for a relationship between short sleep and weight status. Of note, one study of children ages 6–17 years reported no independent association between insufficient sleep and childhood obesity; however, no sleep duration information was provided by parents.⁴⁰

In terms of other metabolic parameters, in a cohort of 652 children, chronic short sleep duration was associated with higher waist circumference, higher insulin levels, and greater HOMA-IR attributed to adiposity.⁴¹ Higher HOMA Indices were also observed in high school-aged children who slept < 6.48 hours.⁴² In children ages 4–7 years, there was increased bone mineral content in children who slept longer than 8 hours.⁴³

A sample of 62 obese 8- to 17-year-old children studied with polysomnography (PSG) showed a U-shape curve for metabolic parameters such as HbA1c and insulin suggesting an ideal range of 7.5–8.5 hours of sleep for this group of children.⁴⁴ A U-shape curve was also observed in a study of 387 children ages 13–17 years, where HOMA-IR was 20% higher when sleep duration was < 5 hours or > 10.5 hours.⁴⁵

Furthermore, higher C-reactive protein was observed in 13–17 year old children with < 9 hours sleep duration.⁴⁶

In summary, short sleep duration is associated with an increased risk for overweight/obesity, particularly in younger children < 10 years, and in boys. There is also evidence that short or long sleep duration is associated with disruption of other metabolic parameters including appetite and glucose/insulin metabolism.

3.4 Mental Health

The variables of interest in the mental health domain related to mood (e.g., depression, anxiety, suicidality, emotional regulation, irritability and self-esteem); risk-taking behaviors (e.g., drug use); problematic behaviors (e.g., defiance and tantrums); and attention-deficit/hyperactivity disorder (ADHD) symptoms (e.g., hyperactivity, impulsivity, and inattentiveness). Studies included approximately 100 to 30,000 participants.

Limited data were available for children in the 4 months to less than 12 months age range. One study looked at the longitudinal sleep trajectories of approximately 3,000 children beginning at age 0–1 year with follow-up at age 6–7 years.⁹ Those children who were persistent short sleepers and poor sleepers had more difficulties with emotional, social and physical functioning at age 6–7 years when compared to “typical sleepers” who slept approximately 14.5 hours of sleep at age 0–1 year and 10.75 hours of sleep at age 6–7 years. Similarly, there were few studies on sleep duration and emotional/mental health within the 12 month to < 3 year old and 3 to < 6 year old age categories. One study collected almost 33,000 parental surveys regarding child sleep duration and the emotional and behavioral regulation of these children at 18 months and 5 years of age.⁴⁷ Children who had short sleep duration of ≤ 10 hours at 18 months were at significantly greater risk of concurrent

and subsequent incident emotional and behavioral problems compared to a reference group of children receiving at least 13 hours of sleep at 18 months.⁴⁷

Studies reviewed for children ages 6–13 years of age focused on associations between total sleep duration and affect, emotional regulation, irritability, relationships with peers/family, and perceived health. Experimental studies of sleep restriction or sleep extension showed that participants sleeping more than 9 hours (mean 9.8 hours) had better emotional lability scores and less restless/impulsive behaviors per teacher reports compared to those sleeping an average of 8.4 hours.⁴⁸ In a similar study, children with a mean of 9.3 hours of sleep had significantly higher positive affect scores and parent-reported emotional regulation compared to children sleeping a mean of 8.1 hours.⁴⁹ In contrast, no group differences were detected in negative affective scores and child-reported emotional regulation ratings. In cross-sectional studies, data were more mixed. For instance, impulsivity scores were significantly higher among children ages 7–8 years sleeping < 7.7 hours compared to those sleeping > 7.7 hours, but no group differences were found in total ADHD global scores.⁵⁰ Likewise, there were no relationships between children sleeping less than 10 hours and their affective scores or reported peer/family relationships, but those sleeping ≥ 10 hours reported better overall health.¹⁵

In contrast, the literature review showed a clearer relationship between sleep duration and mental health among adolescents, 13–18 years of age. Adolescents sleeping ≥ 9 hours of sleep had significantly better life satisfaction scores, fewer general health complaints, and better quality relationships with family compared to those sleeping less.¹⁵ Of greatest concern in the adolescent population are associations between short sleep duration and suicidal thoughts/behaviors and substance abuse. In one cross-sectional survey of 27,939 adolescents, participants who slept 7–8 hours reported less feelings of hopelessness, less tobacco use, less alcohol use, less illicit drug use, fewer suicidal thoughts, and fewer suicidal attempts compared to participants who reported sleeping 6–7 hours per night.⁵¹ Of note, this study showed a negative correlation between more sleep and less concerning behaviors, but after 9 hours of sleep, an increase in these behaviors was noted. This finding suggests a U-shape curve to sleep among adolescents in which too little or too much sleep is associated with mental health problems and substance abuse. To this point, another study showed that teens sleeping ≤ 5 hours and ≥ 10 hours had more suicidal attempts than those sleeping 8 hours per night.⁵²

In summary, there are limited data for infants 4 months to 1 year, but based on the literature, 14.5 hours of sleep seems appropriate. Available longitudinal data on sleep duration for toddlers suggests that at least 13 hours of sleep are beneficial for future mental health outcomes. In school-aged children, there are conflicting data, but children sleeping at least 10 hours report less health complaints and children with < 8 hours of sleep have increased reports of ADHD behaviors. Experimental data suggest at least 9 hours of sleep is necessary for adequate behavioral functioning. In teens, data suggests that 8–9 hours of sleep seems optimal for mental health and < 8 hours is associated with increased inappropriate behavior, including suicidal attempts, and substance abuse. In this population, more than

10 hours of sleep was also associated with an increase in suicidality, although causality cannot be ascertained.

3.5 Immunologic Health

The panel reviewed only 13 studies with respect to loosely defined immunological health and sleep duration. All studies were cross-sectional, observational, or cohort design, and none were randomized controlled studies. The number of participants varied from as few as 54 to just under 1,500. The majority of study participants were older children, adolescents, and teenagers. Sleep duration was measured via a mix of self-report, parent report, actigraphy, or in some cases, the method of measurement was not even reported.

One study suggested that sleep duration of 8–8.9 hours was associated with a healthier immune profile (cortisol, immune cell counts, and cytokines).⁵³ Another study noted an inverse relationship between CRP and sleep duration with higher CRPs noted in those sleeping less than 8 hours.⁴⁶ Other studies showed no relationship between sleep variables and inflammation (fibrinogen, IL-6, and CRP).⁵⁴ No studies suggested too much sleep was harmful.

In summary, based on very limited and less than ideal quality evidence, it would seem that sleep durations less than 8 hours are associated with worse overall immunological health, and that optimal sleep is between 8–9 hours for older children and adolescents.

3.6 Human Performance

The human performance domain included a wide range of outcomes including: neuropsychological testing (e.g., emotional responses, cognitive performance, academic achievement); school grades; rating scales for attention, behavior or executive functioning; ratings scales for behavioral persistence (sustained attention), well-being, health behaviors and complaints; suicide or accident risk; subjective reports of daytime sleepiness; and measured electroencephalography (EEG) arousals. Most studies were cohort or cross-sectional studies of community-based populations; prospective, longitudinal cohort data were rare. Some studies utilized a case-control study design. Other studies used an experimental design to compare the effects of sleep deprivation or sleep extension to the control condition. The number of participants ranged from less than 100 in most of the case-control or experimental studies to over 3,000 in some of the cohort studies.

In a small experimental sample of 10 children aged 30–36 months, loss of a daytime nap resulting in total 24-h sleep duration of less than 11 hours was associated with more negative emotional responses.⁵⁵ In preschoolers, sleep durations of less than 8 hours in the previous 24 hours were associated with a more than 2-fold increased risk of accidental falls.⁵⁶ In a large prospective longitudinal cohort study of 1,492 children aged 6 years, it was found that children with sleep durations less than 10 hours as toddlers and preschoolers had higher parent-reported hyperactivity indices and lower cognitive performance.⁵⁷ In 812 school-aged children, sleeping less than 9 hours per night was associated with worse teacher-reported academic achievement, but did not impact domains of attention, motivation, or relationships.⁵⁸ In 74 children ages 6–12 years,

one week of sleep restriction to 6.5 hours resulted in increases in teacher-reported academic difficulties and attentional problems compared with “optimized” sleep (at least 9 hours).⁵⁹ In 33 children ages 7–11 years with a mean baseline sleep duration of 9 hours 20 minutes, sleep extension of approximately 30 minutes was associated with improved alertness; whereas modest sleep restriction (approximately 1 hour) had opposite effects.⁴⁸ Among 77 children ages 9–12 years with a baseline sleep duration of 8.1 hours, cognitive performance testing improved with a 1-hour extension, but worsened with a 1-hour restriction.⁶⁰ In 32 children ages 8–12 years with a baseline sleep duration of 8.8 hours, 4 days of sleep restriction to 8 hours compared to 4 days of sleep extension to 9.3 hours, resulted in impairment of memory and attention.⁴⁹

Several studies reported on both school-age and teen participants. Self-reported sleep durations of less than 10 hours for children and less than 9 hours for adolescents were associated with increased somatic health complaints (higher odds ratios) when compared to their full sleep conditions.¹⁵ Among 3,011 children 6 to 16 years, short sleep duration (less than the median hours for each of 4 age groups defined by school grade levels) was associated with self-reported daytime tiredness in school.⁶¹ In a large sample of 1,691 participants ages 10–19 years, sleep duration that was one hour less than “optimum” self-reported sleep (which varied by age) predicted lower academic achievement scores, with a larger effect on the older group.⁶² This study also suggested a U-shape effect where much longer sleep durations were associated with worse performance.⁶² In a small sample of 16 adolescents, 5 nights of experimental sleep restriction to 6.5 hours resulted in inattention, diminished learning and lowered arousal in a simulated classroom when compared to a “healthy” sleep condition of 8 hours.⁶³ Lower parent-ratings of executive function and participant performance were associated with self-reported sleepiness in an urban cohort of 236 teens, but not with sleep duration measured by actigraphy.⁶⁴ Among a large cohort of 2,716 teens, sleeping less than 8 hours per night was related to more self-reported tiredness, inferior behavioral persistence (sustained attention), less positive attitude toward life, and lower school grades, as compared to sleeping longer durations.⁶⁵ Daytime tiredness and behavioral persistence mediated the relationship between short sleep duration and positive attitude toward life and school grades. Except for self-reported tiredness, there were no differences between sleeping 8–9 hours and sleeping more than 9 hours. Students who started school 20 minutes later received reliably more sleep and reported less tiredness.⁶⁵ In a large cohort of 1,564 teens, sleeping less than 8 hours was associated with lower cognitive performance for boys, but not girls.⁶⁶ Finally, in a large cohort of 1,362 teens, sleeping less than 8 hours, compared to more than 9 hours, per night was associated with greater risk of suicide, while sleeping 8 to 9 hours was not.⁶⁷

In summary, among various outcomes that comprise human performance, there are sufficient and compelling data to support recommendations for sleep durations of at least 11 hours for toddlers, 10 hours for preschoolers, 9 hours for school-aged children, and 8 hours for teenagers to support optimal performance.

3.7 Developmental Health

Many studies in the developmental health domain targeted the associations between sleep duration and neurobehavioral and neurocognitive measures. There was significant overlap in studies reviewed for this domain, most often with the mental health and human performance domains. Sample sizes ranged from 10 participants (randomized cross-over design) to 35,956 (meta-analysis). There were often methodological issues about assumptions of directionality of relationships and variation in reference categories used for “normal sleep duration.” The covariates differed widely by study and there was significant heterogeneity in outcomes. Results also differed by gender and ethnicity. The panel members noted that besides being associated with child development, short sleep duration has the potential to affect parent-child interactions, which increases the complexity of health interactions.

Short sleep duration has a negative association with cognitive performance in a number of age groups based on longitudinal (< 10 hours,⁵⁷ < 9 hours,⁵⁸ < 11 hours⁶⁸) and cross-sectional (< 9 hours^{15,69}), (< 8 hours⁷⁰), (< 11 hours⁷¹) studies; however, some studies have reported a U-shape relationship with too much sleep time associated with poorer cognitive performance.⁶⁸

For children aged 11–13 months, the proportion of total daily sleep occurring at night has been positively correlated with the development of communication and problem solving skills.⁷² Young children’s poorer language function was positively associated with shorter sleep hours during the night.⁷³ For preschoolers (4 years) and school-aged children (10 years), short sleep duration (< 11 hours) was negatively associated with receptive vocabulary.⁷⁴ Poor emotional regulation and reactivity have been associated with short sleep duration (30–36 month-olds,⁵⁵ 18-month-olds⁴⁷ for < 13 hours, and 2.5–6 year-olds for < 10 hours⁵⁷), but one study reported a U-shape relationship.⁷⁵ Shorter sleep duration has been associated with poorer morning and daytime mood in infants.⁷⁶ As mentioned above in the mental health domain, short sleep duration at 18 months (< 13 hours) predicted more emotionally reactive and aggressive behavior at 5 years.⁴⁷ Persistent short sleep duration (from 0 to 7 years) has been associated with higher levels of child irritability and poorer emotional functioning.⁹

Additionally, naps may be important. Missing one nap by preschoolers produced more negative emotional responses⁵⁵ and nap provision improved toddlers’ performances on a generalization task⁷⁷ and grammatical language patterns 24 hours later⁷⁸ compared with non-nap groups.

In one study, sleeping less than 8 hours per night was associated with poorer cognitive performance on reasoning abilities for adolescent boys but not adolescent girls.⁶⁶ Another study showed that adolescents had positive associations between sleeping less than 8 hours and poorer school grades.⁶⁵ The single meta-analysis of 35,956 children aged 5 to 12 years did not specify sleep duration, but reported that longer sleep duration was associated with better cognitive performance (e.g., implicit memory, intelligence, and school performance) and fewer behavioral problems.⁷⁹ Shorter sleep duration has been linked to school-aged children’s and adolescents’ attention

problems,^{49,59,63,80} restless/impulsive behavior⁴⁸ (< 9 hours) and deterioration in vigilance⁶⁰ (30-minute sleep restriction).

The cited range for normal sleep duration is wide, ranging from 14.5 hours during infancy⁹ to 9–10 hours at school entry and pre-adolescent children^{9,81,82} to about 8.5 hours in late adolescence.⁸³ Studies do not consistently exclude forms of sleep problems. For example, in younger children, the proportion of sleep at night, sleep efficiency and difficulty settling demonstrate the complexity of explicating relationships between sleep duration and developmental well-being.^{72,73} A large, cross-sectional study of 4–9-year-olds did not demonstrate consistent relationships between sleep duration thresholds and learning, based on parent-completed 24-hour time-use diaries.⁸⁴ Only at ages 6 to 7 years did they report a positive relationship between at least 10 hours of sleep per night and mathematical thinking, language, and literacy based on cutoff points on diagnostic categories from teacher report. The findings were difficult to interpret because plots showed most children averaged between 10 to 12 hours of sleep per night. Thus the variance in sleep duration appeared quite limited and the dichotomizing of measures into problem/no problem reduced the information available for testing.

In summary, there appears to be consistent relationships between inadequate sleep duration and development of children's cognition, language, memory, executive function, emotional regulation, and reactivity.

3.8 Longevity

This area focused on sleep patterns in pediatric age groups and how those patterns affect longevity. Seventy-four studies were identified in the initial review. Thirty-seven of the studies associated sleep duration with accidents, suicidal ideation, insomnia, stress, depression, or anxiety. The other 37 studies associated sleep duration with cardiovascular, metabolic, pulmonary, allergy/atopy, and obesity-related conditions. None of the articles either evaluated or demonstrated any direct relationship between sleep duration and longevity in the pediatric population. Although some of the associations, such as increased risk of suicidal ideation or adverse cardiovascular risk factors could impact longevity, those outcomes were not specifically addressed in any of the studies. Therefore, no recommendation was made for optimal sleep duration at any age as related to longevity.

4.0 STRENGTHS AND WEAKNESSES OF THE LITERATURE

The panel recommendations were based on literature characterized by several strengths. Taken together, studies on sleep duration include data on hundreds of thousands of children and teenagers, studied across several continents, aggregated over several decades. The studies include cross-sectional and longitudinal epidemiologic designs, randomized controlled trials, meta-analyses, and a range of other designs. Studies in the mental health category appear to have the strongest evidence.

A number of important limitations in both epidemiologic and lab-based studies are also evident. First, most of the

studies reviewed were cross-sectional, thus findings were correlational, precluding statements of causation. Second, sleep duration was ascertained for a limited time frame around the assessment, with multiple ages grouped together despite obvious differences in development. For example, some studies included participants aged 8–18 years, and it is unknown how the results would have differed had the participants' data been grouped as pre-adolescent vs adolescent with a different age cut-off. Some studies may have had insufficient adjustment for confounders. Additionally, without prospective, long-term studies, pediatric sleep duration data on outcomes such as longevity are lacking.


Furthermore, the methods of sleep assessment present limitations. Most studies described sleep duration obtained from parental report, or child report when age-appropriate. This may be less accurate than averages from daily report on sleep diaries. Parent- or self-reported duration may over- or underestimate sleep duration measured objectively by actigraphy or polysomnography. Additionally, studies varied in how sleep duration was reported, i.e., per 24 hours or overnight sleep. Duration reporting was discussed extensively by panel members to determine the best means to evaluate the available literature and provide appropriate sleep duration recommendations in the predefined age groups of children. Survey questions often had good face validity, but most were not formally validated with psychometric analyses. Multiple surveys also gathered health outcomes such as height, weight, and lists of medical problems, by parental or child report; relying on these reports may be a source of measurement error. Also, measures of sleep duration often do not capture information about the regularity of sleep patterns, the timing of sleep, or the quality of sleep, all of which could directly affect sleep duration and health outcomes.

Experimental designs have important limitations as well. Most studies included short periods of sleep duration manipulation, almost always < 2 weeks. The acute effects of sleep deprivation may not approximate real-world effects of chronic sleep deprivation. Few studies examined sleep duration in the range between 8 and 12 hours. This likely reflects the aim of experimental studies, which often try to maximize differences in outcomes by contrasting extreme sleep duration groups. The absence of experimental groups in these sleep duration ranges creates some uncertainty in recommendations due to lack of data. Many studies also lack generalizability because of small sample sizes that may not represent the population in terms of age, sex, race, ethnicity, socioeconomic status, or health history. Many studies investigated groups of children across multiple ages at different developmental stages, but reported a single result. PSG measurements of sleep duration may not be an accurate measure of typical sleep duration given the different sleep environment (lab) and weak correlations with parental or child self-reported duration.

5.0 VOTING SUMMARY

Voting results from all 3 rounds are presented in **Figures 1–3**. Napping was discussed for age appropriate groups, and reflected in the final recommendations for children through age

Figure 2—Round 2 voting results.



Round 2 voting was based on voting across all age groups under the following seven domains: cardiovascular health, developmental health, human performance, general health, immunology, mental health and metabolic health.

5 years explicitly, with the understanding that some healthy children up to age 7 years may habitually nap. The panel discussed several reasons for why short or long sleep durations may be inappropriate for each age group after incorporating findings from the literature and criteria for sleep disorders. All values are per 24 hours.

For infants, there was consensus that < 12 hours of sleep per 24 hours was inappropriate to support optimal health. There was also consensus that a range of 12 to 16 hours was appropriate to support optimal infant health. Consensus could not be reached on 16 to 18 hours due to uncertainty given the lack of data in this sleep duration range. However, there was consensus that ≥ 18 hours of sleep was inappropriate. There was also consensus that no recommendations could be made in infants < 4 months of age, due to the paucity of published outcomes in infants this young.

For children aged 1 to < 3 years, there was consensus that sleep duration of < 10 hours was inappropriate. There was consensus that 11 to 14 hours per 24 hours was appropriate. Uncertainty was observed in the 10–11 hour range and 14 to 16 hour range, with panel consensus that sleep times longer than 16 hours in this age group were inappropriate.

For children aged 3 to < 6 years, there was consensus that sleep duration of < 10 hours or more than 14 hours was


inappropriate. There was consensus that 10–13 hours of sleep was appropriate. Uncertainty existed in the 13 to 14 hour duration range. The panel discussed that the sleep needs of a 3-year old preschooler may be vastly different than the sleep needs of a 6-year old first-grader.

For children aged 6 years to < 13 years, there was consensus that sleep duration of 9 to 12 hours was appropriate and that > 13 hours was inappropriate. There was uncertainty if 12–13 hours of sleep duration was appropriate.

For teenagers aged 13 to 18 years of age, there was consensus that < 8 hours or > 10 hours of sleep on a regular basis was inappropriate. There was consensus that 8 to 10 hours was appropriate for optimal health.

The panel discussed the merits of recommending sleep durations for multiple age groups vs collapsing all age groups to recommend 1 single sleep duration threshold. Implicit to a range recommendation is the conclusion that sleep duration above a certain amount of sleep is detrimental to health. The literature supported a clear cut-off in optimal sleep duration in the teenager group, as there were unambiguously better health outcomes within the recommended range, and clear negative outcomes and an increased likelihood of sleep disorders when nightly sleep durations were < 8 hours or > 10 hours. Within other age groups, due to the wide age ranges in studies and lack

Figure 3—Round 3 voting results.



Round 3 voting is based on voting summary across all age groups.

of published literature, there was often uncertainty on the outer ends of the agreed upon appropriate sleep duration ranges. The panel also discussed consideration of nap duration for age appropriate children. With these considerations in mind, the panel voted to recommend ranges for each age group. No age groups were collapsed and no age groups were further separated during the in-person meeting because voting in Round 1 had already taken place.

6.0 DISCUSSION

Sleep is a necessity for health. Meeting the need for sleep duration, timing, regularity, and quality requires volitional behaviors by parents and children, which are partially dictated by genetic and physiologic factors. However, a large proportion of inter-individual variability in sleep is likely explained by psychological, behavioral, social, cultural, and environmental factors. Sleep disorders also contribute to this variability.

For reasons stated above, the panel focused solely on the dimension of sleep duration, while recognizing the importance of other factors such as timing, regularity, and quality. The recommendations provide sleep durations that promote optimal pediatric health, but do not address other sleep factors. The panel excluded studies that focused on sleep factors other than sleep duration and studies that focused on total sleep deprivation (no sleep for 24 hours). Total sleep deprivation is not sustainable and thus not reflective of habitual sleep, which was

the focus of the panel. In general, the total sleep time is high in infants and toddlers, with gradual reduction over time as one approaches pre-adolescence and adolescence. This change parallels the ontogeny of cerebral metabolism that was studied in 100 healthy children using positron emission tomography.⁸⁵ The reduction in total sleep time over the first decade is felt to be related to an initial overproduction, and subsequent pruning away of unnecessary synapses.⁸⁵

All panel members agreed that too little sleep for each age group was unhealthy. The panel discussed thoroughly the appropriateness of longer sleep durations for each age group. The panel discussed the higher likelihood of sleep disorders often being associated with longer sleep times, which was then incongruent with sleep duration for *optimal* health. Another consideration was the lack of studies in children on sleep extension. There were a few U-shape curves that showed that both, too little and too much sleep were associated with poorer outcomes. The panel strongly encourages future experimental studies to examine the effects of sleep extension on health outcomes.

The panel recommendations reflect typical or habitual sleep duration, i.e., the ideal average sleep duration. However, the panel discussed that sleep duration in the real world is a dynamic process. Many children, particularly those who have experienced a delay in their circadian rhythms in puberty, have variable sleep schedules depending on school start times, work, and extracurricular activities that may curtail sleep on weekdays and extend sleep on weekends. More studies are needed to better characterize effects of this “social jet lag.”

The panel discussed heterogeneity of sleep duration measurement at length: subjective report vs (more) objective actigraphy, parental report vs child self-report, time in bed vs time asleep, and inclusion vs exclusion of naps. Several studies have shown that self-report may differ substantially from objective measurement.

Polysomnography and actigraphy were utilized in some studies as objective measurements of sleep duration, and the panel discussed the strengths and limitations of these studies. Actigraphy may over-estimate sleep duration relative to self-report in poor sleepers. PSG directly records sleep, however, other factors such as “first night effect” or logistics such as set wake up times to end PSG recordings may be in place and thus not accurately reflect a typical night of sleep. Home sleep apnea testing is not currently recommended in children, and also does not provide accurate reflection of sleep time. As technology improves and the use of activity trackers becomes more widespread in the pediatric population, more data will be available to better measure sleep duration and compare time and effects within large participant samples.

7.0 FUTURE DIRECTIONS

The recommendations of the panel are intended to be a first step towards promoting adequate sleep duration for all children. The panel reviewed existing literature to achieve consensus, and realized that serious gaps in knowledge exist in the pediatric population. As more high-quality literature becomes available, the recommendations may change in the future. Based on this process, specific areas for future research consideration are presented:

1. Implement improved sleep duration measures and study designs, including intervention studies. As child development changes significantly from year to year, it is imperative to study narrower age groups, with validated studies and intervention designs to better delineate sleep deprivation effects on important outcomes such as mental health, physical health, and cognitive functioning. Small preliminary studies have begun to explore the effects of sleep extension on health and functioning in children. More studies are recommended (i.e., between 8 and 10 hours in adolescents) to systematically vary sleep opportunity in discrete steps between the upper and lower amounts using objective measurements of sleep physiology and cognitive performance. These studies should also examine longer time periods (e.g., 30 days), including weekdays and weekend days, to develop more precise dose-response curves for sleep and recovery effects within the ranges most often reported by children and parents. Studies that mimic typical sleep-wake cycles and lifestyles in each age group are needed.
2. Investigate downstream mechanisms linking habitual sleep duration to health and functioning. Intervention studies are needed to help clarify whether modifying sleep duration improves health outcomes in children. Such studies will help to explain whether sleep

duration has a causal role in health and functioning vs operating as a marker of other processes.

3. Delineate the upstream physiologic, behavioral, social, and environmental factors that may play a role in sleep duration and health outcomes. Studies are needed to identify how genetics relate to individual sleep need and the response and resilience to sleep loss. Studies are also needed that explore the roles of race, ethnicity, socioeconomic factors, neighborhoods, school start times, and other factors that contribute to sleep duration.
4. Identify biomarkers of sleep need or sleep deprivation. Inexpensive, reliable, and feasible biomarkers could advance the goals of clinical care, public health, and public policy. In the future, biomarkers may allow clinicians to provide more accurate sleep schedule recommendations to children, parents, and policy makers to facilitate policy decisions (e.g., school start times and teen driving). Biomarkers may greatly assist research assessment of the long term consequences of sleep deprivation.

CONCLUSIONS

The 13-member panel used a modified RAND Appropriateness Method to generate consensus recommendations for the amount of sleep to promote optimal health in children. Multiple rounds of evidence review, discussion and voting were conducted to arrive at the final recommendations. Additional research on the role of sleep duration at different stages in a child's growth and development will help increase the awareness of the importance of sleep and lead to improved health and well-being for children and families.

REFERENCES

1. de Bruin EJ, van Run C, Staaks J, Meijer AM. Effects of sleep manipulation on cognitive functioning of adolescents: a systematic review. *Sleep Med Rev* 2016: in press.
2. Vriend J, Davidson F, Rusak B, Corkum P. emotional and cognitive impact of sleep restriction in children. *Sleep Med Clin* 2015;10:107–15.
3. Cassoff J, Bhatti JA, Gruber R. The effect of sleep restriction on neurobehavioural functioning in normally developing children and adolescents: insights from the attention, behaviour and sleep laboratory. *Pathol Biol (Paris)* 2014;62:319–31.
4. Fitch K, Bernstein S, Aguilar M, et al. The RAND/UCLA Appropriateness Method User's Manual. Santa Monica, CA: RAND, 2001.
5. Paruthi S, Brooks LJ, D'Ambrosio C, et al. Recommended amount of sleep for pediatric populations: a consensus statement of the American Academy of Sleep Medicine. *J Clin Sleep Med* 2016;12:785–6.
6. US Department of Health and Human Services. Sleep Health Objectives. Accessed February 25th, 2016. Available from: <https://www.healthypeople.gov/2020/topics-objectives/topic/sleep-health/objectives>.
7. Oxford Center for Evidence Based Medicine. Accessed March 15, 2016. Available from: <http://www.cebm.net/ocebml-levels-of-evidence/>.
8. Scott N, Blair PS, Emond AM, et al. Sleep patterns in children with ADHD: a population-based cohort study from birth to 11 years. *J Sleep Res* 2013;22:121–8.

9. Magee CA, Gordon R, Caputi P. Distinct developmental trends in sleep duration during early childhood. *Pediatrics* 2014;133:e1561-7.
10. Valent F, Brusafiero S, Barbone F. A case-crossover study of sleep and childhood injury. *Pediatrics* 2001;107:E23.
11. Wang H, Sekine M, Chen X, Yamagami T, Kagamimori S. Lifestyle at 3 years of age and quality of life (QOL) in first-year junior high school students in Japan: results of the Toyama Birth Cohort Study. *Qual Life Res* 2008;17:257-65.
12. Quan SF, Goodwin JL, Babar SI, et al. Sleep architecture in normal Caucasian and Hispanic children aged 6-11 years recorded during unattended home polysomnography: experience from the Tucson Children's Assessment of Sleep Apnea Study (TuCASA). *Sleep Med* 2003;4:13-9.
13. Williams RL, Karacan I, Hirsch CJ. *Electroencephalography (EEG) of human sleep: clinical applications*. New York: John Wiley & Sons, 1974.
14. Olds T, Blunden S, Petkov J, Forchino F. The relationships between sex, age, geography and time in bed in adolescents: a meta-analysis of data from 23 countries. *Sleep Med Rev* 2010;14:371-8.
15. Segura-Jimenez V, Carbonell-Baeza A, Keating XD, Ruiz JR, Castro-Pinero J. Association of sleep patterns with psychological positive health and health complaints in children and adolescents. *Qual Life Res* 2015;24:885-95.
16. Lam LT, Yang L. Short duration of sleep and unintentional injuries among adolescents in China. *Am J Epidemiol* 2007;166:1053-8.
17. Milewski MD, Skaggs DL, Bishop GA, et al. Chronic lack of sleep is associated with increased sports injuries in adolescent athletes. *J Pediatr Orthop* 2014;34:129-33.
18. Sampei M, Murata K, Dakeishi M, Wood DC. Cardiac autonomic hypofunction in preschool children with short nocturnal sleep. *Tohoku J Exp Med* 2006;208:235-42.
19. Guo X, Zheng L, Li Y, et al. Association between sleep duration and hypertension among Chinese children and adolescents. *Clin Cardiol* 2011;34:774-81.
20. Azadbakht L, Kelishadi R, Saraf-Bank S, et al. The association of sleep duration and cardiometabolic risk factors in a national sample of children and adolescents: the CASPIAN III Study. *Nutrition* 2013;29:1133-41.
21. Kuciene R, Dulskiene V. Associations of short sleep duration with prehypertension and hypertension among Lithuanian children and adolescents: a cross-sectional study. *BMC Public Health* 2014;14:255.
22. Lee JA, Park HS. Relation between sleep duration, overweight, and metabolic syndrome in Korean adolescents. *Nutr Metab Cardiovasc Dis* 2014;24:65-71.
23. Paciencia I, Barros H, Araújo J, Ramos E. Association between sleep duration and blood pressure in adolescents. *Hypertens Res* 2013;36:747-52.
24. Berentzen NE, Smit HA, Bekkers MB, et al. Time in bed, sleep quality and associations with cardiometabolic markers in children: the Prevention and Incidence of Asthma and Mite Allergy birth cohort study. *J Sleep Res* 2014;23:3-12.
25. Rey-Lopez JP, de Carvalho HB, de Moraes AC, et al. Sleep time and cardiovascular risk factors in adolescents: the HELENA (Healthy Lifestyle in Europe by Nutrition in Adolescence) study. *Sleep Med* 2014;15:104-10.
26. Sung V, Beebe DW, Vandyke R, et al. Does sleep duration predict metabolic risk in obese adolescents attending tertiary services? A cross-sectional study. *Sleep* 2011;34:891-8.
27. Fatima Y, Doi SA, Mamun AA. Longitudinal impact of sleep on overweight and obesity in children and adolescents: a systematic review and bias-adjusted meta-analysis. *Obes Rev* 2015;16:137-49.
28. Chen X, Beydoun MA, Wang Y. Is sleep duration associated with childhood obesity? A systematic review and meta-analysis. *Obesity (Silver Spring)* 2008;16:265-74.
29. Taveras EM, Rifas-Shiman SL, Oken E, Gunderson EP, Gillman MW. Short sleep duration in infancy and risk of childhood overweight. *Arch Pediatr Adolesc Med* 2008;162:305-11.
30. Bell JF, Zimmerman FJ. Shortened nighttime sleep duration in early life and subsequent childhood obesity. *Arch Pediatr Adolesc Med* 2010;164:840-5.
31. Diethelm K, Bolzenius K, Cheng G, Remer T, Buyken AE. Longitudinal associations between reported sleep duration in early childhood and the development of body mass index, fat mass index and fat free mass index until age 7. *Int J Pediatr Obes* 2011;6:e114-23.
32. Hiscock H, Sczalzo K, Canterford L, Wake M. Sleep duration and body mass index in 0-7-year olds. *Arch Dis Child* 2011;96:735-9.
33. Duncan JS, Schofield G, Duncan EK, Rush EC. Risk factors for excess body fatness in New Zealand children. *Asia Pac J Clin Nutr* 2008;17:138-47.
34. Sekine M, Yamagami T, Handa K, et al. A dose-response relationship between short sleeping hours and childhood obesity: results of the Toyama Birth Cohort Study. *Child Care Health Dev* 2002;28:163-70.
35. Meng LP, Liu AL, Hu X, et al. Report on childhood obesity in China (10): association of sleep duration with obesity. *Biomed Environ Sci* 2012;25:133-40.
36. Drescher AA, Goodwin JL, Silva GE, Quan SF. Caffeine and screen time in adolescence: associations with short sleep and obesity. *J Clin Sleep Med* 2011;7:337-42.
37. Eisenmann JC, Ekkekakis P, Holmes M. Sleep duration and overweight among Australian children and adolescents. *Acta Paediatr* 2006;95:956-63.
38. Garaulet M, Ortega FB, Ruiz JR, et al. Short sleep duration is associated with increased obesity markers in European adolescents: effect of physical activity and dietary habits. The HELENA study. *Int J Obes (Lond)* 2011;35:1308-17.
39. Fisher A, McDonald L, van Jaarsveld CH, et al. Sleep and energy intake in early childhood. *Int J Obes (Lond)* 2014;38:926-9.
40. Hassan F, Davis MM, Chervin RD. No independent association between insufficient sleep and childhood obesity in the National Survey of Children's Health. *J Clin Sleep Med* 2011;7:153-7.
41. Cespedes EM, Rifas-Shiman SL, Redline S, Gillman MW, Peña MM, Taveras EM. Longitudinal associations of sleep curtailment with metabolic risk in mid-childhood. *Obesity (Silver Spring)* 2014;22:2586-92.
42. Matthews KA, Dahl RE, Owens JF, Lee L, Hall M. Sleep duration and insulin resistance in healthy black and white adolescents. *Sleep* 2012;35:1353-8.
43. Casazza K, Hanks LJ, Fernandez JR. Shorter sleep may be a risk factor for impaired bone mass accrual in childhood. *J Clin Densitom* 2011;14:453-7.
44. Koren D, Levitt Katz LE, Brar PC, Gallagher PR, Berkowitz RI, Brooks LJ. Sleep architecture and glucose and insulin homeostasis in obese adolescents. *Diabetes Care* 2011;34:2442-7.
45. Javaheri S, Storer-Isser A, Rosen CL, Redline S. Association of short and long sleep durations with insulin sensitivity in adolescents. *J Pediatr* 2011;158:617-23.
46. Martinez-Gomez D, Eisenmann JC, Gomez-Martinez S, et al.; AFINOS Study Group. Sleep duration and emerging cardiometabolic risk markers in adolescents. The AFINOS study. *Sleep Med* 2011;12:997-1002.
47. Sivertsen B, Harvey AG, Reichborn-Kjennerud T, Torgersen L, Ystrom E, Hysing M. Later emotional and behavioral problems associated with sleep problems in toddlers: a longitudinal study. *JAMA Pediatr* 2015;169:575-82.
48. Gruber R, Cassoff J, Frenette S, Wiebe S, Carrier J. Impact of sleep extension and restriction on children's emotional lability and impulsivity. *Pediatrics* 2012;130:e1155-61.
49. Vriend JL, Davidson FD, Corkum PV, Rusak B, Chambers CT, McLaughlin EN. Manipulating sleep duration alters emotional functioning and cognitive performance in children. *J Pediatr Psychol* 2013;38:1058-69.
50. Paavonen EJ, Raikkonen K, Lahti J, et al. Short sleep duration and behavioral symptoms of attention-deficit/hyperactivity disorder in healthy 7- to 8-year-old children. *Pediatrics* 2009;123:e857-64.
51. Winsler A, Deutsch A, Vorona RD, Payne PA, Szklo-Coxe M. Sleepless in Fairfax: the difference one more hour of sleep can make for teen hopelessness, suicidal ideation, and substance use. *J Youth Adolesc* 2015;44:362-78.
52. Fitzgerald CT, Messias E, Buysse DJ. Teen sleep and suicidality: results from the youth risk behavior surveys of 2007 and 2009. *J Clin Sleep Med* 2011;7:351-6.
53. Perez de Heredia, Garaulet M, Gómez-Martínez S, et al. Self-reported sleep duration, white blood cell counts and cytokine profiles in European adolescents: the HELENA study. *Sleep Med* 2014;15:1251-8.
54. Countryman AJ, Saab PG, Llabre MM, Penedo FJ, McCalla JR, Schneiderman N. Cardiometabolic risk in adolescents: associations with physical activity, fitness, and sleep. *Ann Behav Med* 2013;45:121-31.
55. Berger RH, Miller AL, Seifer R, Cares SR, LeBourgeois MK. Acute sleep restriction effects on emotion responses in 30- to 36-month-old children. *J Sleep Res* 2012;21:235-46.

56. Boto LR, Crispim JN, de Melo IS, et al. Sleep deprivation and accidental fall risk in children. *Sleep Med* 2012;13:88–95.
57. Touchette E, Petit D, Seguin JR, Boivin M, Tremblay RE, Montplaisir JY. Associations between sleep duration patterns and behavioral/cognitive functioning at school entry. *Sleep* 2007;30:1213–9.
58. Li S, Arguelles L, Jiang F, et al. Sleep, school performance, and a school-based intervention among school-aged children: a sleep series study in China. *PLoS One* 2013;8:e67928.
59. Fallone G, Acebo C, Seifer R, Carskadon MA. Experimental restriction of sleep opportunity in children: effects on teacher ratings. *Sleep* 2005;28:1561–7.
60. Sadeh A, Gruber R, Raviv A. The effects of sleep restriction and extension on school-age children: what a difference an hour makes. *Child Dev* 2003;74:444–55.
61. Garmy P, Nyberg P, Jakobsson U. Sleep and television and computer habits of Swedish school-age children. *J Sch Nurs* 2012;28:469–76.
62. Eide E. Sleep and Student Achievement. *Eastern Econ J* 2012;38.
63. Beebe DW, Rose D, Amin R. Attention, learning, and arousal of experimentally sleep-restricted adolescents in a simulated classroom. *J Adolesc Health* 2010;47:523–5.
64. Anderson B, Storfer-Isser A, Taylor HG, Rosen CL, Redline S. Associations of executive function with sleepiness and sleep duration in adolescents. *Pediatrics* 2009;123:e701–7.
65. Perkinson-Gloor N, Lemola S, Grob A. Sleep duration, positive attitude toward life, and academic achievement: the role of daytime tiredness, behavioral persistence, and school start times. *J Adolesc* 2013;36:311–8.
66. Ortega FB, Ruiz JR, Castillo R, et al. Sleep duration and cognitive performance in adolescence. The AVENA study. *Acta Paediatr* 2010;99:454–6.
67. Liu X. Sleep and adolescent suicidal behavior. *Sleep* 2004;27:1351–8.
68. Kocovska D, Rijlaarsdam J, Ghassabian A, et al. Early childhood sleep patterns and cognitive development at age 6 years: the Generation R Study. *J Pediatric Psychol* 2016;1–9.
69. Ravid S, Afek I, Suraiya S, Shahar E, Pillar G. Kindergarten children's failure to qualify for first grade could result from sleep disturbances. *J Child Neurol* 2009;24:816–22.
70. Stea TH, Knutsen T, Torstveit MK. Association between short time in bed, health-risk behaviors and poor academic achievement among Norwegian adolescents. *Sleep Med* 2014;15:666–71.
71. Tso W, Rao N, Jiang F, et al. Sleep duration and school readiness of Chinese preschool children. *J Pediatr* 2016;169:266–71.
72. Gibson R, Elder A, Gander P. Actigraphic sleep and developmental progress of one-year-old infants. *Sleep Biol Rhythms* 2012;10:77–83.
73. Iemura A, Iwasaki M, Yamakawa N, et al. Influence of sleep-onset time on the development of 18-month-old infants: Japan Children's cohort study. *Brain Dev* 2016;38:364–72.
74. Seegers V, Touchette E, Dionne G, et al. Short persistent sleep duration is associated with poor receptive vocabulary performance in middle childhood. *J Sleep Res* 2016;25:325–32.
75. Carvalho Bos S, Gomes A, Clemente V, et al. Sleep and behavioral/emotional problems in children: a population-based study. *Sleep Med* 2009;10:66–74.
76. Mindell J, Lee C. Sleep, mood, and development in infants. *Infant Behav Dev* 2015;41:102–7.
77. Horváth K, Liu S, Plunkett K. A daytime nap facilitates generalization of word meanings in young toddlers. *Sleep* 2016;39:203–7.
78. Hupbach A, Gomez RL, Bootzin RR, Nadel L. Nap-dependent learning in infants. *Dev Sci* 2009;12:1007–12.
79. Astill RG, Van der Heijden KB, Van IJendoorn MH, Van Someren EJW. Sleep, cognition, and behavioral problems in school-age children: a century of research meta-analyzed. *Psychol Bull* 2012;138:1109–38.
80. Lo JC, Ong JL, Leong RLF, Gooley JJ, Chee MWL. Cognitive performance, sleepiness, and mood in partially sleep deprived adolescents: the need for sleep study. *Sleep* 2016;39:687–98.
81. Lemola S, Raikkonen K, Scheier MF, et al. Sleep quantity, quality, and optimism in children. *J Sleep Res* 2011;20:12–20.
82. Jiang X, Hardy LL, Bauer LA, et al. Sleep duration, schedule and quality among urban Chinese children and adolescents: associations with routine after-school activities. *PLoS One* 2015;10:e0115326.
83. Olds T, Maher C, Blunden S, Matricciani L. Normative data on the sleep habits of Australian children and adolescents. *Sleep* 2010;33:1381–8.
84. Price AM, Quach J, Wake M, Bittman M, Hiscock H. Cross-sectional sleep thresholds for optimal health and well-being in Australian 4 to 9-year-olds. *Sleep Med* 2016;22:83–90.
85. Chugani HT, Phelps ME, Mazziotta JC. Positron emission tomography study of human brain functional development. *Ann Neurol* 1987;22:487–97.

ACKNOWLEDGMENTS

The AASM acknowledges the following individuals for their participation and contributions: Non-Participating Observers: Michael Twery, PhD, National Health, Lung, Blood Institute, NIH, Bethesda, MD; and Joel Porquez, RST, RPSGT, CCSH, American Association of Sleep Technologists. American Academy of Sleep Medicine Staff: John Noel; Jonathan L. Heald, MA; and Sarah Hashmi, MBBS, MSc, MPH.

SUBMISSION & CORRESPONDENCE INFORMATION

Submitted for publication September, 2016

Accepted for publication September, 2016

Address correspondence to: Shalini Paruthi, MD, St. Luke's Hospital, 232 S. Woods Mill Road, St. Louis, MO 63017; Email: research@aasmnet.org

DISCLOSURE STATEMENT

Funding for this project was provided by the American Academy of Sleep Medicine, and supported by the cooperative agreement number 1U50DP004930-03 from the Centers for Disease Control and Prevention (CDC). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the CDC. Dr. Maski has consulted for Medscape Inc. and has received research support from Jazz Pharmaceuticals. Dr. Rosen has consulted as a medical advisor for Jazz Pharmaceuticals. The other Consensus Conference Panel members have indicated no financial conflicts of interest.

LITERATURE SEARCH TERMS

Category	Search Term		
Sleep Duration	Napping	[All fields]	
	Sleep debt	[All fields]	
	Short sleep	[All fields]	
	Long sleep	[All fields]	
	Sleep deprivation	[All fields]	
	Insufficient sleep syndrome	[All fields]	
	Excessive sleep	[All fields]	
	Sleep curtailment	[All fields]	
	Sleep length	[All fields]	
	Sleep need	[All fields]	
	Sleep duration	[All fields]	
	Sleep deficiency	[All fields]	
	Cancer	Neoplasms	[MeSH terms]
	Cardiovascular Health	Cardiovascular diseases	[MeSH terms]
Heart valves		[MeSH terms]	
Heart valve diseases		[MeSH terms]	
Blood pressure		[MeSH terms]	
Developmental Health	Child Development Disorders, Pervasive	[MeSH terms]	
	Academic performance	[All fields]	
	Behavior	[MeSH terms]	
	Psychology, educational	[MeSH terms]	
	Personality	[MeSH terms]	
	Learning	[MeSH terms]	
	Attention	[MeSH terms]	
	School start time	[All fields]	
	Neuropsychological tests	[MeSH terms]	
	Neuropsycholo*	[All fields]	
	Neurocognitive	[All fields]	
	Neurobehavioral	[All fields]	
	Emotional regulation	[All fields]	
	Language	[MeSH terms]	
Socialization	[MeSH terms]		

Category	Search Term	
Human Performance	Function	[All fields]
	Activities of daily living	[MeSH terms]
	Alertness	[All fields]
	Cognition	[MeSH terms]
	Patient safety	[All fields]
	Patient harms	[All fields]
	Accidents	[MeSH terms]
	Accident proneness	[MeSH terms]
	Fatigue	[MeSH terms]
	Vigilance	[All fields]
	Psychomotor performance	[MeSH terms]
	Reaction	[All fields]
	Comprehension	[MeSH terms]
	Confusion	[MeSH terms]
	Impairment	[All fields]
	Sleepiness <i>NOT sleep disorders</i>	[All fields]
	Sleep satisfaction	[All fields]
	Achievement	[MeSH terms]
	Athletic performance	[MeSH terms]
	Automobile driving	[MeSH terms]

Category	Search Term	
General Health	Epidemiology	[MeSH terms]
	Prevention medicine	[MeSH terms]
	Emergencies	[All fields]
	Endemic diseases	[All fields]
	Environmental medicine	[All fields]
	Environmental microbiology	[MeSH terms]
	Disease transmission, infectious	[MeSH terms]
	Environmental pollution	[MeSH terms]
	Public health practice	[MeSH terms]
	Wellness	[All fields]
	Well-being	[All fields]
	Quality of life	[MeSH terms]
	Wounds and Injuries	[MeSH terms]
	Immunological Health	Inflammation
Autoimmune disease		[MeSH terms]
Cytokines		[MeSH terms]
Leukocytes		[MeSH terms]
Phagocytes		[MeSH terms]
Immunoproteins		[MeSH terms]
Vaccination		[MeSH terms]
Asthma		[MeSH terms]
Common cold		[MeSH terms]
Infection		[MeSH terms]
Longevity		Death
	Survival	[MeSH terms]
	Vital statistics	[MeSH terms]
	Suicide	[MeSH terms]

Category	Search Term		
Mental Health	Mental Health	[MeSH terms]	
	Depression	[MeSH terms]	
	Psychiatry	[MeSH terms]	
	Anxiety disorders	[MeSH terms]	
	Mood disorders	[MeSH terms]	
	Schizophrenia and disorders with psychotic features	[MeSH terms]	
	Substance-related disorders	[MeSH terms]	
	Mental Disorders Diagnosed in Childhood	[MeSH terms]	
	Attention Deficit and Disruptive Behavior Disorders	[MeSH terms]	
	Eating disorders	[MeSH terms]	
	Emotions	[MeSH terms]	
	Metabolic Health	Metabolism	[MeSH terms]
		Body weight	[MeSH terms]
Metabolic disease		[MeSH terms]	
Insulin		[MeSH terms]	
Orexins		[MeSH terms]	
Ghrelin		[MeSH terms]	
Adipokines		[MeSH terms]	
Waist-hip ratio		[MeSH terms]	
Thyroid gland		[MeSH terms]	
Pediatric obesity		[MeSH terms]	
Pain		Pain	[MeSH terms]
	Nociception	[MeSH terms]	
	Rheumatic diseases	[MeSH terms]	

REFERENCE LIST OF FINAL EVIDENCE

Cardiovascular Health

- Au CT, Ho CK, Wing YK, Lam HS, Li AM. Acute and chronic effects of sleep duration on blood pressure. *Pediatrics*. 2014 Jan;133(1):e64-72.
- Azadbakht L, Kelishadi R, Khodarahmi M, Qorbani M, Heshmat R, Motlagh ME, Taslimi M, Ardalan G. The association of sleep duration and cardiometabolic risk factors in a national sample of children and adolescents: the CASPIAN III study. *Nutrition*. 2013 Sep;29(9):1133-41.
- Berentzen NE, Smit HA, Bekkers MB, Brunekreef B, Koppelman GH, De Jongste JC, Kerkhof M, Van Rossem L, Wijga AH. Time in bed, sleep quality and associations with cardiometabolic markers in children: the Prevention and Incidence of Asthma and Mite Allergy birth cohort study. *J Sleep Res*. 2014 Feb;23(1):3-12.
- Countryman AJ, Saab PG, Llabre MM, Penedo FJ, McCalla JR, Schneiderman N. Cardiometabolic risk in adolescents: associations with physical activity, fitness, and sleep. *Ann Behav Med*. 2013 Feb;45(1):121-31.
- Guo X, Zheng L, Li Y, Yu S, Liu S, Zhou X, Zhang X, Sun Z, Wang R, Sun Y. Association between sleep duration and hypertension among Chinese children and adolescents. *Clin Cardiol*. 2011 Dec;34(12):774-81.
- Javaheri S, Storfer-Isser A, Rosen CL, Redline S. Sleep quality and elevated blood pressure in adolescents. *Circulation*. 2008 Sep 2;118(10):1034-40.
- Kuciene R, Dulskiene V. Associations of short sleep duration with prehypertension and hypertension among Lithuanian children and adolescents: a cross-sectional study. *BMC Public Health*. 2014 Mar 15;14:255.
- Lee JA, Park HS. Relation between sleep duration, overweight, and metabolic syndrome in Korean adolescents. *Nutr Metab Cardiovasc Dis*. 2014 Jan;24(1):65-71.
- Lehto R, Ray C, Lahti-Koski M, Roos E. Health behaviors, waist circumference and waist-to-height ratio in children. *Eur J Clin Nutr*. 2011 Jul;65(7):841-8.
- Mahler B, Kamperis K, Schroeder M, Frøkiær J, Djurhuus JC, Rittig S. Sleep deprivation induces excess diuresis and natriuresis in healthy children. *Am J Physiol Renal Physiol*. 2012 Jan 15;302(2):F236-43.
- Martikainen S, Pesonen AK, Feldt K, Jones A, Lahti J, Pyhälä R, Heinonen K, Kajantie E, Eriksson J, Räikkönen K. Poor sleep and cardiovascular function in children. *Hypertension*. 2011 Jul;58(1):16-21.
- Martinez-Gomez D, Eisenmann JC, Gomez-Martinez S, Hill EE, Zapatera B, Veiga OL, Marcos A; AFINOS Study Group. Sleep duration and emerging cardiometabolic risk markers in adolescents. The AFINOS study. *Sleep Med*. 2011 Dec;12(10):997-1002.
- Meininger JC, Gallagher MR, Eissa MA, Nguyen TQ, Chan W. Sleep duration and its association with ambulatory blood pressure in a school-based, diverse sample of adolescents. *Am J Hypertens*. 2014 Jul;27(7):948-55.
- Paciência I, Barros H, Araújo J, Ramos E. Association between sleep duration and blood pressure in adolescents. *Hypertens Res*. 2013 Aug;36(8):747-52.
- Peach H, Gaultney JF, Reeve CL. Sleep characteristics, body mass index, and risk for hypertension in young adolescents. *J Youth Adolesc*. 2015 Feb;44(2):271-84.
- Rey-López JP, de Carvalho HB, de Moraes AC, Ruiz JR, Sjöström M, Marcos A, Polito A, Gottrand F, Manios Y, Kafatos A, Molnar D, Widhalm K, De Henauw S, Moreno LA; HELENA study. Sleep time and cardiovascular risk factors in adolescents: the HELENA (Healthy Lifestyle in Europe by Nutrition in Adolescence) study. *Sleep Med*. 2014 Jan;15(1):104-10.
- Sampei M, Murata K, Dakeishi M, Wood DC. Cardiac autonomic hypofunction in preschool children with short nocturnal sleep. *Tohoku J Exp Med*. 2006 Mar;208(3):235-42.
- Sampei M, Dakeishi M, Wood DC, Murata K. Impact of total sleep duration on blood pressure in preschool children. *Biomed Res*. 2006 Jun;27(3):111-5.
- Sung V, Beebe DW, Vandyke R, Fenchel MC, Crimmins NA, Kirk S, Hiscock H, Amin R, Wake M. Does sleep duration predict metabolic risk in obese adolescents attending tertiary services? A cross-sectional study. *Sleep*. 2011 Jul 1;34(7):891-8.
- Wells JC, Hallal PC, Reichert FF, Menezes AM, Araújo CL, Victora CG. Sleep patterns and television viewing in relation to obesity and blood pressure: evidence from an adolescent Brazilian birth cohort. *Int J Obes (Lond)*. 2008 Jul;32(7):1042-9.

Developmental Health

- Abe T, Hagihara A, Nobutomo K. Sleep patterns and impulse control among Japanese junior high school students. *J Adolesc*. 2010 Oct;33(5):633-41.
- Adolescent Sleep Working Group; Committee on Adolescence; Council on School Health. School start times for adolescents. *Pediatrics*. 2014 Sep;134(3):642-9.
- Agras WS, Hammer LD, McNicholas F, Kraemer HC. Risk factors for childhood overweight: a prospective study from birth to 9.5 years. *J Pediatr*. 2004 Jul;145(1):20-5. Erratum in: *J Pediatr*. 2004 Sep;145(3):424.
- Aishworiya R, Chan P, Kiing J, Chong SC, Laino AG, Tay SKh. Sleep behaviour in a sample of preschool children in Singapore. *Ann Acad Med Singapore*. 2012 Mar;41(3):99-104.
- Al-Hazzaa HM, Musaiger AO, Abahussain NA, Al-Sobayel HI, Qahwaji DM. Lifestyle correlates of self-reported sleep duration among Saudi adolescents: a multicentre school-based cross-sectional study. *Child Care Health Dev*. 2014 Jul;40(4):533-42.
- Anderson B, Storfer-Isser A, Taylor HG, Rosen CL, Redline S. Associations of executive function with sleepiness and sleep duration in adolescents. *Pediatrics*. 2009 Apr;123(4):e701-7.
- Androutsos O, Moschonis G, Mavrogianni C, Roma-Giannikou E, Chrousos GP, Kanaka-Gantenbein C, Manios Y. Identification of lifestyle patterns, including sleep deprivation, associated with insulin resistance in children: the Healthy Growth Study. *Eur J Clin Nutr*. 2014 Mar;68(3):344-9.

- Anuntaseree W, Mo-suwan L, Vasiknanonte P, Kuasirikul S, Ma-a-lee A, Choprapawan C. Night waking in Thai infants at 3 months of age: association between parental practices and infant sleep. *Sleep Med.* 2008 Jul;9(5):564-71.
- Appelhans BM, Fitzpatrick SL, Li H, Cail V, Waring ME, Schneider KL, Whited MC, Busch AM, Pagoto SL. The home environment and childhood obesity in low-income households: indirect effects via sleep duration and screen time. *BMC Public Health.* 2014 Nov 9;14:1160.
- Arakawa M, Taira K, Tanaka H, Yamakawa K, Toguchi H, Kadekaru H, Yamamoto Y, Uezu E, Shirakawa S. A survey of junior high school students' sleep habit and lifestyle in Okinawa. *Psychiatry Clin Neurosci.* 2001 Jun;55(3):211-2.
- Aronen ET, Paavonen EJ, Fällberg M, Soininen M, Törrönen J. Sleep and psychiatric symptoms in school-age children. *J Am Acad Child Adolesc Psychiatry.* 2000 Apr;39(4):502-8.
- Arora T, Taheri S. Associations among late chronotype, body mass index and dietary behaviors in young adolescents. *Int J Obes (Lond).* 2015 Jan;39(1):39-44.
- Arora T, Hosseini-Araghi M, Bishop J, Yao GL, Thomas GN, Taheri S. The complexity of obesity in U.K. adolescents: relationships with quantity and type of technology, sleep duration and quality, academic performance and aspiration. *Pediatr Obes.* 2013 Oct;8(5):358-66.
- Asaka Y, Takada S. Activity-based assessment of the sleep behaviors of VLBW preterm infants and full-term infants at around 12 months of age. *Brain Dev.* 2010 Feb;32(2):150-5.
- Asarnow LD, McGlinchey E, Harvey AG. The effects of bedtime and sleep duration on academic and emotional outcomes in a nationally representative sample of adolescents. *J Adolesc Health.* 2014 Mar;54(3):350-6.
- Astill RG, Van der Heijden KB, Van Ijzendoorn MH, Van Someren EJ. Sleep, cognition, and behavioral problems in school-age children: a century of research meta-analyzed. *Psychol Bull.* 2012 Nov;138(6):1109-38.
- BaHammam A, Bin Saeed A, Al-Faris E, Shaikh S. Sleep duration and its correlates in a sample of Saudi elementary school children. *Singapore Med J.* 2006 Oct;47(10):875-81.
- Ball GD, Lenk JM, Barbarich BN, Plotnikoff RC, Fishburne GJ, Mackenzie KA, Willows ND. Overweight children and adolescents referred for weight management: are they meeting lifestyle behaviour recommendations? *Appl Physiol Nutr Metab.* 2008 Oct;33(5):936-45.
- Barnes JC, Meldrum RC. The impact of sleep duration on adolescent development: a genetically informed analysis of identical twin pairs. *J Youth Adolesc.* 2015 Feb;44(2):489-506.
- Baum KT, Desai A, Field J, Miller LE, Rausch J, Beebe DW. Sleep restriction worsens mood and emotion regulation in adolescents. *J Child Psychol Psychiatry.* 2014;55(2):180-90.
- Beebe DW, Simon S, Summer S, Hemmer S, Strotman D, Dolan LM. Dietary intake following experimentally restricted sleep in adolescents. *Sleep.* 2013 Jun 1;36(6):827-34.
- Beebe DW, Ris MD, Kramer ME, Long E, Amin R. The association between sleep disordered breathing, academic grades, and cognitive and behavioral functioning among overweight subjects during middle to late childhood. *Sleep.* 2010 Nov;33(11):1447-56.
- Beebe DW, Rose D, Amin R. Attention, learning, and arousal of experimentally sleep-restricted adolescents in a simulated classroom. *J Adolesc Health.* 2010 Nov;47(5):523-5.
- Beebe DW, Fallone G, Godiwala N, Flanigan M, Martin D, Schaffner L, Amin R. Feasibility and behavioral effects of an at-home multi-night sleep restriction protocol for adolescents. *J Child Psychol Psychiatry.* 2008 Sep;49(9):915-23.
- Bel S, Michels N, De Vriendt T, Patterson E, Cuenca-García M, Diethelm K, Gutin B, Grammatikaki E, Manios Y, Leclercq C, Ortega FB, Moreno LA, Gottrand F, Gonzalez-Gross M, Widhalm K, Kafatos A, Garaulet M, Molnar D, Kaufman JM, Gilbert CC, Hallström L, Sjöström M, et al. Association between self-reported sleep duration and dietary quality in European adolescents. *Br J Nutr.* 2013 Sep 14;110(5):949-59.
- Berger RH, Miller AL, Seifer R, Cares SR, LeBourgeois MK. Acute sleep restriction effects on emotion responses in 30- to 36-month-old children. *J Sleep Res.* 2012 Jun;21(3):235-46.
- Bertelle V, Sevestre A, Laou-Hap K, Nagahapitiye MC, Sizun J. Sleep in the neonatal intensive care unit. *J Perinat Neonatal Nurs.* 2007 Apr-Jun;21(2):140-8; quiz 149-50. Review.
- Bertelle V, Mabin D, Adrien J, Sizun J. Sleep of preterm neonates under developmental care or regular environmental conditions. *Early Hum Dev.* 2005 Jul;81(7):595-600.
- Bibiloni Mdel M, Martinez E, Llull R, Juarez MD, Pons A, Tur JA. Prevalence and risk factors for obesity in Balearic Islands adolescents. *Br J Nutr.* 2010 Jan;103(1):99-106.
- Biggs SN, Lushington K, van den Heuvel CJ, Martin AJ, Kennedy JD. Inconsistent sleep schedules and daytime behavioral difficulties in school-aged children. *Sleep Med.* 2011 Sep;12(8):780-6.
- Bruni O, Baumgartner E, Sette S, Ancona M, Caso G, Di Cosimo ME, Mannini A, Ometto M, Pasquini A, Ulliana A, Ferri R. Longitudinal study of sleep behavior in normal infants during the first year of life. *J Clin Sleep Med.* 2014 Oct 15;10(10):1119-27.
- Bruni O, Russo PM, Ferri R, Novelli L, Galli F, Guidetti V. Relationships between headache and sleep in a non-clinical population of children and adolescents. *Sleep Med.* 2008 Jul;9(5):542-8.
- Buckhalt JA, El-Sheikh M, Keller P. Children's sleep and cognitive functioning: race and socioeconomic status as moderators of effects. *Child Dev.* 2007 Jan-Feb;78(1):213-31.
- Burt J, Dube L, Thibault L, Gruber R. Sleep and eating in childhood: a potential behavioral mechanism underlying the relationship between poor sleep and obesity. *Sleep Med.* 2014 Jan;15(1):71-5.

- Calamaro CJ, Park S, Mason TB, Marcus CL, Weaver TE, Pack A, Ratcliffe SJ. Shortened sleep duration does not predict obesity in adolescents. *J Sleep Res.* 2010 Dec;19(4):559-66
- Calamaro CJ, Mason TB, Ratcliffe SJ. Adolescents living the 24/7 lifestyle: effects of caffeine and technology on sleep duration and daytime functioning. *Pediatrics.* 2009 Jun;123(6):e1005-10.
- Capaldi Li VF, Handwerker K, Richardson E, Stroud LR. Associations between sleep and cortisol responses to stress in children and adolescents: a pilot study. *Behav Sleep Med.* 2005;3(4):177-92.
- Carskadon MA. Sleep's effects on cognition and learning in adolescence. *Prog Brain Res.* 2011;190:137-43.
- Carskadon MA, Wolfson AR, Acebo C, Tzischinsky O, Seifer R. Adolescent sleep patterns, circadian timing, and sleepiness at a transition to early school days. *Sleep.* 1998 Dec 15;21(8):871-81.
- Carvalho Bos S, Gomes A, Clemente V, Marques M, Pereira AT, Maia B, Soares MJ, Cabral AS, Macedo A, Gozal D, Azevedo MH. Sleep and behavioral/emotional problems in children: a population-based study. *Sleep Med.* 2009 Jan;10(1):66-74.
- Casoff J, Bhatti JA, Gruber R. The effect of sleep restriction on neurobehavioural functioning in normally developing children and adolescents: insights from the Attention, Behaviour and Sleep Laboratory. *Pathol Biol (Paris).* 2014 Oct;62(5):319-31.
- Chung KF, Cheung MM. Sleep-wake patterns and sleep disturbance among Hong Kong Chinese adolescents. *Sleep.* 2008 Feb;31(2):185-94.
- Cicek B, Ozturk A, Mazicioglu MM, Elmali F, Turp N, Kurtoglu S. The risk analysis of arm fat area in Turkish children and adolescents. *Ann Hum Biol.* 2009 Jan-Feb;36(1):28-37.
- Cortesi F, Giannotti F, Sebastiani T, Vagnoni C, Marioni P. Cosleeping versus solitary sleeping in children with bedtime problems: child emotional problems and parental distress. *Behav Sleep Med.* 2008;6(2):89-105
- Countryman AJ, Saab PG, Llabre MM, Penedo FJ, McCalla JR, Schneiderman N. Cardiometabolic risk in adolescents: associations with physical activity, fitness, and sleep. *Ann Behav Med.* 2013 Feb;45(1):121-31.
- Crosby B, LeBourgeois MK, Harsh J. Racial differences in reported napping and nocturnal sleep in 2- to 8-year-old children. *Pediatrics.* 2005 Jan;115(1 Suppl):225-32.
- Dahl RE. Biological, developmental, and neurobehavioral factors relevant to adolescent driving risks. *Am J Prev Med.* 2008 Sep;35(3 Suppl):S278-84.
- Dahl RE, Lewin DS. Pathways to adolescent health sleep regulation and behavior. *J Adolesc Health.* 2002 Dec;31(6 Suppl):175-84. Review.
- Davis AL, Avis KT, Schwebel DC. The effects of acute sleep restriction on adolescents' pedestrian safety in a virtual environment. *J Adolesc Health.* 2013 Dec;53(6):785-90.
- de Jong E, Stocks T, Visscher TL, HiraSing RA, Seidell JC, Renders CM. Association between sleep duration and overweight: the importance of parenting. *Int J Obes (Lond).* 2012 Oct;36(10):1278-84
- Dev DA, McBride BA, Fiese BH, Jones BL, Cho H; Behalf Of The Strong Kids Research Team. Risk factors for overweight/obesity in preschool children: an ecological approach. *Child Obes.* 2013 Oct;9(5):399-408.
- Dewald JF, Short MA, Gradisar M, Oort FJ, Meijer AM. The Chronic Sleep Reduction Questionnaire (CSRQ): a cross-cultural comparison and validation in Dutch and Australian adolescents. *J Sleep Res.* 2012 Oct;21(5):584-94.
- Dexter D, Bijwadia J, Schilling D, Applebaugh G. Sleep, sleepiness and school start times: a preliminary study. *WMJ.* 2003;102(1):44-6.
- Do YK, Shin E, Bautista MA, Foo K. The associations between self-reported sleep duration and adolescent health outcomes: what is the role of time spent on Internet use? *Sleep Med.* 2013 Feb;14(2):195-200.
- Doane LD, Thurston EC. Associations among sleep, daily experiences, and loneliness in adolescence: evidence of moderating and bidirectional pathways. *J Adolesc.* 2014 Feb;37(2):145-54.
- Dollman J, Ridley K, Olds T, Lowe E. Trends in the duration of school-day sleep among 10- to 15-year-old South Australians between 1985 and 2004. *Acta Paediatr.* 2007 Jul;96(7):1011-4.
- Dorofaeff TF, Denny S. Sleep and adolescence. Do New Zealand teenagers get enough? *J Paediatr Child Health.* 2006 Sep;42(9):515-20.
- Duncan JS, Schofield G, Duncan EK, Rush EC. Risk factors for excess body fatness in New Zealand children. *Asia Pac J Clin Nutr.* 2008;17(1):138-47.
- Eaton DK, McKnight-Eily LR, Lowry R, Perry GS, Presley-Cantrell L, Croft JB. Prevalence of insufficient, borderline, and optimal hours of sleep among high school students - United States, 2007. *J Adolesc Health.* 2010 Apr;46(4):399-401.
- El-Sheikh M, Buckhalt JA, Mark Cummings E, Keller P. Sleep disruptions and emotional insecurity are pathways of risk for children. *J Child Psychol Psychiatry.* 2007 Jan;48(1):88-96.
- Eisenburg LK, Corpeleijn E, van Sluijs EM, Atkin AJ. Clustering and correlates of multiple health behaviours in 9-10 year old children. *PLoS One.* 2014;9(6):e99498.
- Ertan P, Yilmaz O, Caglayan M, Sogut A, Aslan S, Yuksel H. Relationship of sleep quality and quality of life in children with monosymptomatic enuresis. *Child Care Health Dev.* 2009 Jul;35(4):469-74.
- Escribano C, Díaz-Morales JF. Daily fluctuations in attention at school considering starting time and chronotype: an exploratory study. *Chronobiol Int.* 2014 Jul;31(6):761-9.
- Fallone G, Acebo C, Seifer R, Carskadon MA. Experimental restriction of sleep opportunity in children: effects on teacher ratings. *Sleep.* 2005 Dec;28(12):1561-7.
- Fallone G, Acebo C, Arnedt JT, Seifer R, Carskadon MA. Effects of acute sleep restriction on behavior, sustained attention, and response inhibition in children. *Percept Mot Skills.* 2001 Aug;93(1):213-29.
- Firouzi S, Poh BK, Ismail MN, Sadeghilar A. Sleep habits, food intake, and physical activity levels in normal and overweight and obese Malaysian children. *Obes Res Clin Pract.* 2014 Jan-Feb;8(1):e70-8.

- Fitzgerald CT, Messias E, Buysse DJ. Teen sleep and suicidality: results from the youth risk behavior surveys of 2007 and 2009. *J Clin Sleep Med*. 2011 Aug 15;7(4):351-6.
- Fredriksen K, Rhodes J, Reddy R, Way N. Sleepless in Chicago: tracking the effects of adolescent sleep loss during the middle school years. *Child Dev*. 2004 Jan-Feb;75(1):84-95.
- Fukuda K, Ishihara K. Routine evening naps and night-time sleep patterns in junior high and high school students. *Psychiatry Clin Neurosci*. 2002 Jun;56(3):229-30.
- Gaina A, Sekine M, Hamanishi S, Chen X, Wang H, Yamagami T, Kagamimori S. Daytime sleepiness and associated factors in Japanese school children. *J Pediatr*. 2007 Nov;151(5):518-22, 522.e1-4.
- Garaulet M, Ortega FB, Ruiz JR, Rey-López JP, Béghin L, Manios Y, Cuenca-García M, Plada M, Diethelm K, Kafatos A, Molnár D, Al-Tahan J, Moreno LA. Short sleep duration is associated with increased obesity markers in European adolescents: effect of physical activity and dietary habits. The HELENA study. *Int J Obes (Lond)*. 2011 Oct;35(10):1308-17.
- Garmy P, Nyberg P, Jakobsson U. Sleep and television and computer habits of Swedish school-age children. *J Sch Nurs*. 2012 Dec;28(6):469-76.
- Gau SS, Shang CY, Merikangas KR, Chiu YN, Soong WT, Cheng AT. Association between morningness-eveningness and behavioral/emotional problems among adolescents. *J Biol Rhythms*. 2007 Jun;22(3):268-74.
- Geiger A, Achermann P, Jenni OG. Association between sleep duration and intelligence scores in healthy children. *Dev Psychol*. 2010 Jul;46(4):949-54.
- Giannotti F, Cortesi F, Cerquiglini A, Miraglia D, Vagnoni C, Sebastiani T, Bernabei P. An investigation of sleep characteristics, EEG abnormalities and epilepsy in developmentally regressed and non-regressed children with autism. *J Autism Dev Disord*. 2008 Nov;38(10):1888-97.
- Gibson ES, Powles AC, Thabane L, O'Brien S, Molnar DS, Trajanovic N, Ogilvie R, Shapiro C, Yan M, Chilcott-Tanser L. "Sleepiness" is serious in adolescence: two surveys of 3235 Canadian students." *BMC Public Health*. 2006 May 2;6:116.
- Golley RK, Maher CA, Matricciani L, Olds TS. Sleep duration or bedtime? Exploring the association between sleep timing behaviour, diet and BMI in children and adolescents. *Int J Obes (Lond)*. 2013 Apr;37(4):546-51.
- Gruber R, Cassoff J, Frenette S, Wiebe S, Carrier J. Impact of sleep extension and restriction on children's emotional lability and impulsivity. *Pediatrics*. 2012 Nov;130(5):e1155-61.
- Gruber R, Wiebe S, Montecalvo L, Brunetti B, Amsel R, Carrier J. Impact of sleep restriction on neurobehavioral functioning of children with attention deficit hyperactivity disorder. *Sleep*. 2011 Mar 1;34(3):315-23.
- Gruber R, Laviolette R, Deluca P, Monson E, Cornish K, Carrier J. Short sleep duration is associated with poor performance on IQ measures in healthy school-age children. *Sleep Med*. 2010 Mar;11(3):289-94.
- Guo X, Zheng L, Li Y, Yu S, Sun G, Yang H, Zhou X, Zhang X, Sun Z, Sun Y. Differences in lifestyle behaviors, dietary habits, and familial factors among normal-weight, overweight, and obese Chinese children and adolescents. *Int J Behav Nutr Phys Act*. 2012 Oct 2;9:120.
- Gössel-Sybank R, Grimmer I, Korte J, Siegmund R. Actigraphic monitoring of the activity-rest behavior of preterm and full-term infants at 20 months of age. *Chronobiol Int*. 2004 Jul;21(4-5):661-71.
- Hagenauer MH, Perryman JI, Lee TM, Carskadon MA. Adolescent changes in the homeostatic and circadian regulation of sleep. *Dev Neurosci*. 2009;31(4):276-84.
- Haines J, McDonald J, O'Brien A, Sherry B, Bottino CJ, Schmidt ME, Taveras EM. Healthy Habits, Happy Homes: randomized trial to improve household routines for obesity prevention among preschool-aged children. *JAMA Pediatr*. 2013 Nov;167(11):1072-9.
- Harrington SA. Relationships of objectively measured physical activity and sleep with BMI and academic outcomes in 8-year-old children. *Appl Nurs Res*. 2013 May;26(2):63-70.
- Hart CN, Carskadon MA, Considine RV, Fava JL, Lawton J, Raynor HA, Jelalian E, Owens J, Wing R. Changes in children's sleep duration on food intake, weight, and leptin. *Pediatrics*. 2013 Dec;132(6):e1473-80.
- Hiscock H, Sciberras E, Mensah F, Gerner B, Efron D, Khano S, Oberklaid F. Impact of a behavioural sleep intervention on symptoms and sleep in children with attention deficit hyperactivity disorder, and parental mental health: randomised controlled trial. *BMJ*. 2015 Jan 20;350:h68.
- Hjorth MF, Chaput JP, Ritz C, Dalskov SM, Andersen R, Astrup A, Tetens I, Michaelsen KF, Sjödin A. Fatness predicts decreased physical activity and increased sedentary time, but not vice versa: support from a longitudinal study in 8- to 11-year-old children. *Int J Obes (Lond)*. 2014 Jul;38(7):959-65.
- Hjorth MF, Quist JS, Andersen R, Michaelsen KF, Tetens I, Astrup A, Chaput JP, Sjödin A. Change in sleep duration and proposed dietary risk factors for obesity in Danish school children. *Pediatr Obes*. 2014 Dec;9(6):e156-9.
- Hodge D, Carollo TM, Lewin M, Hoffman CD, Sweeney DP. Sleep patterns in children with and without autism spectrum disorders: developmental comparisons. *Res Dev Disabil*. 2014 Jul;35(7):1631-8.
- Holley S, Hill CM, Stevenson J. An hour less sleep is a risk factor for childhood conduct problems. *Child Care Health Dev*. 2011 Jul;37(4):563-70.
- Holm SM, Forbes EE, Ryan ND, Phillips ML, Tarr JA, Dahl RE. Reward-related brain function and sleep in pre/early pubertal and mid/late pubertal adolescents. *J Adolesc Health*. 2009 Oct;45(4):326-34.
- Honomichl RD, Goodlin-Jones BL, Burnham M, Gaylor E, Anders TF. Sleep patterns of children with pervasive developmental disorders. *J Autism Dev Disord*. 2002 Dec;32(6):553-61.
- Humphreys JS, Gringras P, Blair PS, Scott N, Henderson J, Fleming PJ, Emond AM. Sleep patterns in children with autistic spectrum disorders: a prospective cohort study. *Arch Dis Child*. 2014 Feb;99(2):114-8.

- Hupbach A, Gomez RL, Bootzin RR, Nadel L. Nap-dependent learning in infants. *Dev Sci*. 2009 Nov;12(6):1007-12.
- Hysing M, Harvey AG, Torgersen L, Ystrom E, Reichborn-Kjennerud T, Sivertsen B. Trajectories and predictors of nocturnal awakenings and sleep duration in infants. *J Dev Behav Pediatr*. 2014 Jun;35(5):309-16.
- Ikeda M, Kaneita Y, Kondo S, Itani O, Ohida T. Epidemiological study of sleep habits among four-and-a-half-year-old children in Japan. *Sleep Med*. 2012 Aug;13(7):787-94.
- Jang SI, Lee KS, Park EC. Relationship between current sleep duration and past suicidal ideation or attempt among Korean adolescents. *J Prev Med Public Health*. 2013 Nov;46(6):329-35.
- Jarrin DC, McGrath JJ, Drake CL. Beyond sleep duration: distinct sleep dimensions are associated with obesity in children and adolescents. *Int J Obes (Lond)*. 2013 Apr;37(4):552-8.
- Javaheri S, Storfer-Isser A, Rosen CL, Redline S. Association of short and long sleep durations with insulin sensitivity in adolescents. *J Pediatr*. 2011 Apr;158(4):617-23.
- Jiang X, Hardy LL, Baur LA, Ding D, Wang L, Shi H. Sleep duration, schedule and quality among urban Chinese children and adolescents: associations with routine after-school activities. *PLoS One*. 2015;10(1):e0115326.
- Kang SG, Lee YJ, Kim SJ, Lim W, Lee HJ, Park YM, Cho IH, Cho SJ, Hong JP. Weekend catch-up sleep is independently associated with suicide attempts and self-injury in Korean adolescents. *Compr Psychiatry*. 2014 Feb;55(2):319-25.
- Keller PS, El-Sheikh M, Buckhalt JA. Children's attachment to parents and their academic functioning: sleep disruptions as moderators of effects. *J Dev Behav Pediatr*. 2008 Dec;29(6):441-9.
- Kelly RJ, El-Sheikh M. Reciprocal relations between children's sleep and their adjustment over time. *Dev Psychol*. 2014 Apr;50(4):1137-47.
- Kim HS, Ham OK, Kim JW, Park JY. Association between sleep duration and psychological health in overweight and obese children in Korea. *Nurs Health Sci*. 2012 Jun;14(2):238-43.
- Kirjavainen J, Lehtonen L, Kirjavainen T, Kero P; 24-Hour Ambulatory Sleep Polygraphy study. Sleep of excessively crying infants: a 24-Hour Ambulatory Sleep Polygraphy study. *Pediatrics*. 2004 Sep;114(3):592-600.
- Kjeldsen JS, Hjorth MF, Andersen R, Michaelsen KF, Tetens I, Astrup A, Chaput JP, Sjödin A. Short sleep duration and large variability in sleep duration are independently associated with dietary risk factors for obesity in Danish school children. *Int J Obes (Lond)*. 2014 Jan;38(1):32-9.
- Kleiser C, Schaffrath Rosario A, Mensink GB, Prinz-Langenohl R, Kurth BM. Potential determinants of obesity among children and adolescents in Germany: results from the cross-sectional KiGGS Study. *BMC Public Health*. 2009 Feb 2;9:46.
- Klingenberg L, Chaput JP, Holmbäck U, Jennum P, Astrup A, Sjödin A. Sleep restriction is not associated with a positive energy balance in adolescent boys. *Am J Clin Nutr*. 2012 Aug;96(2):240-8.
- Kohyama J, Shiiki T, Ohinata-Sugimoto J, Hasegawa T. Potentially harmful sleep habits of 3-year-old children in Japan. *J Dev Behav Pediatr*. 2002 Apr;23(2):67-70.
- Komada Y, Abe T, Okajima I, Asaoka S, Matsuura N, Usui A, Shirakawa S, Inoue Y. Short sleep duration and irregular bedtime are associated with increased behavioral problems among Japanese preschool-age children. *Tohoku J Exp Med*. 2011;224(2):127-36.
- Kopasz M, Loessl B, Valerius G, Koenig E, Matthaeas N, Hornyak M, Kloepfer C, Nissen C, Riemann D, Voderholzer U. No persisting effect of partial sleep curtailment on cognitive performance and declarative memory recall in adolescents. *J Sleep Res*. 2010 Mar;19(1 Pt 1):71-9.
- Kopasz M, Loessl B, Hornyak M, Riemann D, Nissen C, Piosczyk H, Voderholzer U. Sleep and memory in healthy children and adolescents - a critical review. *Sleep Med Rev*. 2010 Jun;14(3):167-77.
- Kostyun RO, Milewski MD, Hafeez I. Sleep disturbance and neurocognitive function during the recovery from a sport-related concussion in adolescents. *Am J Sports Med*. 2015 Mar;43(3):633-40.
- Kouros CD, El-Sheikh M. Daily mood and sleep: reciprocal relations and links with adjustment problems. *J Sleep Res*. 2015 Feb;24(1):24-31.
- Kubiszewski V, Fontaine R, Potard C, Gimenes G. Bullying, sleep/wake patterns and subjective sleep disorders: findings from a cross-sectional survey. *Chronobiol Int*. 2014 May;31(4):542-53.
- Kuciene R, Dulskiene V. Associations of short sleep duration with prehypertension and hypertension among Lithuanian children and adolescents: a cross-sectional study. *BMC Public Health*. 2014 Mar 15;14:255.
- Kuriyan R, Thomas T, Sumithra S, Lokesh DP, Sheth NR, Joy R, Bhat S, Kurpad AV. Potential factors related to waist circumference in urban South Indian children. *Indian Pediatr*. 2012 Feb;49(2):124-8.
- Kuriyan R, Bhat S, Thomas T, Vaz M, Kurpad AV. Television viewing and sleep are associated with overweight among urban and semi-urban South Indian children. *Nutr J*. 2007 Sep 20;6:25.
- Kwon JA, Lee M, Yoo KB, Park EC. Does the duration and time of sleep increase the risk of allergic rhinitis? Results of the 6-year nationwide Korea youth risk behavior web-based survey. *PLoS One*. 2013;8(8):e72507.
- Lam JC, Mahone EM, Mason T, Scharf SM. The effects of napping on cognitive function in preschoolers. *J Dev Behav Pediatr*. 2011 Feb-Mar;32(2):90-7.
- Lazaratou H, Dikeos DG, Anagnostopoulos DC, Sbokou O, Soldatos CR. Sleep problems in adolescence. A study of senior high school students in Greece. *Eur Child Adolesc Psychiatry*. 2005 Jul;14(4):237-43.
- Lee YJ, Cho SJ, Cho IH, Kim SJ. Insufficient sleep and suicidality in adolescents. *Sleep*. 2012 Apr 1;35(4):455-60.
- Lehto JE, Uusitalo-Malmivaara L. Sleep-related factors: associations with poor attention and depressive symptoms. *Child Care Health Dev*. 2014 May;40(3):419-25.

- Lehto R, Ray C, Lahti-Koski M, Roos E. Health behaviors, waist circumference and waist-to-height ratio in children. *Eur J Clin Nutr.* 2011 Jul;65(7):841-8.
- Lemola S, Schwarz B, Siffert A. Interparental conflict and early adolescents' aggression: is irregular sleep a vulnerability factor? *J Adolesc.* 2012 Feb;35(1):97-105..
- Lemola S, Rääkkönen K, Scheier MF, Matthews KA, Pesonen AK, Heinonen K, Lahti J, Komsu N, Paavonen JE, Kajantie E. Sleep quantity, quality and optimism in children. *J Sleep Res.* 2011 Mar;20(1 Pt 1):12-20.
- Lewandowski AS, Palermo TM, De la Motte S, Fu R. Temporal daily associations between pain and sleep in adolescents with chronic pain versus healthy adolescents. *Pain.* 2010 Oct;151(1):220-5.
- Li S, Arguelles L, Jiang F, Chen W, Jin X, Yan C, Tian Y, Hong X, Qian C, Zhang J, Wang X, Shen X. Sleep, school performance, and a school-based intervention among school-aged children: a sleep series study in China. *PLoS One.* 2013 Jul 10;8(7):e67928.
- Li Y, Jin H, Owens JA, Hu C. The association between sleep and injury among school-aged children in rural China: a case-control study. *Sleep Med.* 2008 Jan;9(2):142-8.
- Lin WH, Yi CC. Unhealthy sleep practices, conduct problems, and daytime functioning during adolescence. *J Youth Adolesc.* 2015 Feb;44(2):431-46.
- Liu X, Hubbard JA, Fabes RA, Adam JB. Sleep disturbances and correlates of children with autism spectrum disorders. *Child Psychiatry Hum Dev.* 2006 Winter;37(2):179-91.
- Liu X. Sleep and adolescent suicidal behavior. *Sleep.* 2004 Nov 1;27(7):1351-8.
- Liu X, Uchiyama M, Okawa M, Kurita H. Prevalence and correlates of self-reported sleep problems among Chinese adolescents. *Sleep.* 2000 Feb 1;23(1):27-34.
- Loessl B, Valerius G, Kopasz M, Hornyak M, Riemann D, Voderholzer U. Are adolescents chronically sleep-deprived? An investigation of sleep habits of adolescents in the Southwest of Germany. *Child Care Health Dev.* 2008 Sep;34(5):549-56.
- Lytle LA, Pasch KE, Farbakhsh K. The relationship between sleep and weight in a sample of adolescents. *Obesity (Silver Spring).* 2011 Feb;19(2):324-31.
- Magee CA, Gordon R, Caputi P. Distinct developmental trends in sleep duration during early childhood. *Pediatrics.* 2014 Jun;133(6):e1561-7.
- Mak KK, Lee SL, Ho SY, Lo WS, Lam TH. Sleep and academic performance in Hong Kong adolescents. *J Sch Health.* 2012 Nov;82(11):522-7.
- Malow BA, Marzec ML, McGrew SG, Wang L, Henderson LM, Stone WL. Characterizing sleep in children with autism spectrum disorders: a multidimensional approach. *Sleep.* 2006 Dec;29(12):1563-71.
- Martinez-Gomez D, Eisenmann JC, Gomez-Martinez S, Hill EE, Zapatera B, Veiga OL, Marcos A; AFINOS Study Group. Sleep duration and emerging cardiometabolic risk markers in adolescents. The AFINOS study. *Sleep Med.* 2011 Dec;12(10):997-1002.
- Martinez-Gomez D, Moreno LA, Romeo J, Rey-López P, Castillo R, Cabero MJ, Vicente-Rodriguez G, Gutiérrez A, Veiga OL. Combined influence of lifestyle risk factors on body fat in Spanish adolescents--the Avena study. *Obes Facts.* 2011;4(2):105-11.
- Maski KP, Kothare SV. Sleep deprivation and neurobehavioral functioning in children. *Int J Psychophysiol.* 2013 Aug;89(2):259-64.
- Matamura M, Tochigi M, Usami S, Yonehara H, Fukushima M, Nishida A, Togo F, Sasaki T. Associations between sleep habits and mental health status and suicidality in a longitudinal survey of monozygotic twin adolescents. *J Sleep Res.* 2014 Jun;23(3):290-4.
- Matsushita M, Koyama A, Ushijima H, Mikami A, Katsumata Y, Kikuchi Y, Ichimi N, Jono T, Fujise N, Ikeda M. Sleep duration and its association with sleepiness and depression in ronin-sei" preparatory school students." *Asian J Psychiatr.* 2014 Jun;9:61-6.
- McDermott BM, Mamun AA, Najman JM, Williams GM, O'callaghan MJ, Bor W. Preschool children perceived by mothers as irregular eaters: physical and psychosocial predictors from a birth cohort study. *J Dev Behav Pediatr.* 2008 Jun;29(3):197-205.
- Meneilly J, Tremblay MS, Leduc G, Boyer C, Bélanger P, Leblanc AG, Borghese MM, Chaput JP. Objectively-measured sleep and its association with adiposity and physical activity in a sample of Canadian children. *J Sleep Res.* 2015 Apr;24(2):131-9.
- Meijer AM. Chronic sleep reduction, functioning at school and school achievement in preadolescents. *J Sleep Res.* 2008 Dec;17(4):395-405. Erratum in: *J Sleep Res.* 2009 Mar;18(1):144.
- Meininger JC, Gallagher MR, Eissa MA, Nguyen TQ, Chan W. Sleep duration and its association with ambulatory blood pressure in a school-based, diverse sample of adolescents. *Am J Hypertens.* 2014 Jul;27(7):948-55.
- Meldrum RC, Barnes JC, Hay C. Sleep deprivation, low self-control, and delinquency: a test of the strength model of self-control. *J Youth Adolesc.* 2015 Feb;44(2):465-77.
- Mercer PW, Merritt SL, Cowell JM. Differences in reported sleep need among adolescents. *J Adolesc Health.* 1998 Nov;23(5):259-63.
- Michels N, Clays E, De Buyzere M, Vanaelst B, De Henauw S, Sioen I. Children's sleep and autonomic function: low sleep quality has an impact on heart rate variability. *Sleep.* 2013 Dec 1;36(12):1939-46.
- Molfese DL, Ivanenko A, Key AF, Roman A, Molfese VJ, O'Brien LM, Gozal D, Kota S, Hudac CM. A one-hour sleep restriction impacts brain processing in young children across tasks: evidence from event-related potentials. *Dev Neuropsychol.* 2013;38(5):317-36.
- Monasta L, Batty GD, Cattaneo A, Lutje V, Ronfani L, Van Lenthe FJ, Brug J. Early-life determinants of overweight and obesity: a review of systematic reviews. *Obes Rev.* 2010 Oct;11(10):695-708.
- Montgomery-Downs HE, Gozal D. Sleep habits and risk factors for sleep-disordered breathing in infants and young toddlers in Louisville, Kentucky. *Sleep Med.* 2006 Apr;7(3):211-9.
- Moore M, Kirchner HL, Drotar D, Johnson N, Rosen C, Redline S. Correlates of adolescent sleep time and variability in sleep time: the role of individual and health

- related characteristics. *Sleep Med.* 2011 Mar;12(3):239-45.
- Moore M, Kirchner HL, Drotar D, Johnson N, Rosen C, Ancoli-Israel S, Redline S. Relationships among sleepiness, sleep time, and psychological functioning in adolescents. *J Pediatr Psychol.* 2009 Nov-Dec;34(10):1175-83.
- Morley BC, Scully ML, Niven PH, Okely AD, Baur LA, Pratt IS, Wakefield MA; NaSSDA Study Team. What factors are associated with excess body weight in Australian secondary school students? *Med J Aust.* 2012 Feb 20;196(3):189-92.
- Moseley L, Gradisar M. Evaluation of a school-based intervention for adolescent sleep problems. *Sleep.* 2009 Mar;32(3):334-41.
- Must A, Parisi SM. Sedentary behavior and sleep: paradoxical effects in association with childhood obesity. *Int J Obes (Lond).* 2009 Apr;33 Suppl 1:S82-6.
- Natal CL, Lourenço TJ, Silva LA, Boscolo RA, Silva A, Tufik S, Mello MT. Gender differences in the sleep habits of 11-13 year olds. *Rev Bras Psiquiatr.* 2009 Dec;31(4):358-61.
- Ng EP, Ng DK, Chan CH. Sleep duration, wake/sleep symptoms, and academic performance in Hong Kong Secondary School Children. *Sleep Breath.* 2009 Nov;13(4):357-67.
- Nixon GM, Thompson JM, Han DY, Becroft DM, Clark PM, Robinson E, Waldie KE, Wild CJ, Black PN, Mitchell EA. Short sleep duration in middle childhood: risk factors and consequences. *Sleep.* 2008 Jan;31(1):71-8.
- Noland H, Price JH, Dake J, Telljohann SK. Adolescents' sleep behaviors and perceptions of sleep. *J Sch Health.* 2009 May;79(5):224-30.
- O'Brien EM, Mindell JA. Sleep and risk-taking behavior in adolescents. *Behav Sleep Med.* 2005;3(3):113-33.
- O'Dea JA, Dibley MJ, Rankin NM. Low sleep and low socioeconomic status predict high body mass index: a 4-year longitudinal study of Australian schoolchildren. *Pediatr Obes.* 2012 Aug;7(4):295-303.
- Ohida T, Osaki Y, Doi Y, Tanihata T, Minowa M, Suzuki K, Wada K, Suzuki K, Kaneita Y. An epidemiologic study of self-reported sleep problems among Japanese adolescents. *Sleep.* 2004 Aug 1;27(5):978-85.
- Olds T, Maher C, Blunden S, Matricciani L. Normative data on the sleep habits of Australian children and adolescents. *Sleep.* 2010 Oct;33(10):1381-8.
- Olds T, Blunden S, Petkov J, Forchino F. The relationships between sex, age, geography and time in bed in adolescents: a meta-analysis of data from 23 countries. *Sleep Med Rev.* 2010 Dec;14(6):371-8.
- Olds TS, Maher CA, Matricciani L. Sleep duration or bedtime? Exploring the relationship between sleep habits and weight status and activity patterns. *Sleep.* 2011 Oct 1;34(10):1299-307.
- Ortega FB, Ruiz JR, Labayen I, Kwak L, Harro J, Oja L, Veidebaum T, Sjöström M. Sleep duration and activity levels in Estonian and Swedish children and adolescents. *Eur J Appl Physiol.* 2011 Oct;111(10):2615-23.
- Ortega FB, Chillón P, Ruiz JR, Delgado M, Albers U, Alvarez-Granda JL, Marcos A, Moreno LA, Castillo MJ. Sleep patterns in Spanish adolescents: associations with TV watching and leisure-time physical activity. *Eur J Appl Physiol.* 2010 Oct;110(3):563-73.
- Owens J. Classification and epidemiology of childhood sleep disorders. *Prim Care.* 2008 Sep;35(3):533-46, vii.
- Owens J, Maxim R, McGuinn M, Nobile C, Msall M, Alario A. Television-viewing habits and sleep disturbance in school children. *Pediatrics.* 1999 Sep;104(3):e27.
- Owens JA, Belon K, Moss P. Impact of delaying school start time on adolescent sleep, mood, and behavior. *Arch Pediatr Adolesc Med.* 2010 Jul;164(7):608-14.
- Owens JA, Fernando S, Mc Guinn M. Sleep disturbance and injury risk in young children. *Behav Sleep Med.* 2005;3(1):18-31.
- Paavonen EJ, Räikkönen K, Pesonen AK, Lahti J, Komsu N, Heinonen K, Järvenpää AL, Strandberg T, Kajantie E, Porkka-Heiskanen T. Sleep quality and cognitive performance in 8-year-old children. *Sleep Med.* 2010 Apr;11(4):386-92.
- Paavonen EJ, Vehkalahti K, Vanhala R, von Wendt L, Nieminen-von Wendt T, Aronen ET. Sleep in children with Asperger syndrome. *J Autism Dev Disord.* 2008 Jan;38(1):41-51.
- Paavonen EJ, Nieminen-von Wendt T, Vanhala R, Aronen ET, von Wendt L. Effectiveness of melatonin in the treatment of sleep disturbances in children with Asperger disorder. *J Child Adolesc Psychopharmacol.* 2003 Spring;13(1):83-95.
- Paciência I, Barros H, Araújo J, Ramos E. Association between sleep duration and blood pressure in adolescents. *Hypertens Res.* 2013 Aug;36(8):747-52.
- Paksarian D, Rudolph KE, He JP, Merikangas KR. School Start Time and Adolescent Sleep Patterns: Results From the U.S. National Comorbidity Survey--Adolescent Supplement. *Am J Public Health.* 2015 Jul;105(7):1351-7.
- Paraskakis E, Ntouros T, Ntokos M, Siavana O, Bitsori M, Galanakis E. Siesta and sleep patterns in a sample of adolescents in Greece. *Pediatr Int.* 2008 Oct;50(5):690-3.
- Park JH, Yoo JH, Kim SH. Associations between non-restorative sleep, short sleep duration and suicidality: findings from a representative sample of Korean adolescents. *Psychiatry Clin Neurosci.* 2013 Jan;67(1):28-34.
- Park S. The association between short sleep duration and body mass index among South Korean children and adolescents. *J Sch Nurs.* 2013 Apr;29(2):142-50.
- Park YM, Matsumoto K, Seo YJ, Shinkoda H. Sleep and chronotype for children in Japan. *Percept Mot Skills.* 1999 Jun;88(3 Pt 2):1315-29.
- Pasch KE, Latimer LA, Cance JD, Moe SG, Lytle LA. Longitudinal bi-directional relationships between sleep and youth substance use. *J Youth Adolesc.* 2012 Sep;41(9):1184-96.
- Pasch KE, Laska MN, Lytle LA, Moe SG. Adolescent sleep, risk behaviors, and depressive symptoms: are they linked? *Am J Health Behav.* 2010 Mar-Apr;34(2):237-48.
- Peach H, Gaultney JF, Reeve CL. Sleep characteristics, body mass index, and risk for hypertension in young adolescents. *J Youth Adolesc.* 2015 Feb;44(2):271-84.
- Perez-Chada D, Perez-Lloret S, Videla AJ, Cardinali D, Bergna MA, Fernández-Acquier M, Larrateguy L, Zabert

- GE, Drake C. Sleep disordered breathing and daytime sleepiness are associated with poor academic performance in teenagers. A study using the Pediatric Daytime Sleepiness Scale (PDSS). *Sleep*. 2007 Dec;30(12):1698-703.
- Perez-Lloret S, Videla AJ, Richaudeau A, Vigo D, Rossi M, Cardinali DP, Perez-Chada D. A multi-step pathway connecting short sleep duration to daytime somnolence, reduced attention, and poor academic performance: an exploratory cross-sectional study in teenagers. *J Clin Sleep Med*. 2013 May 15;9(5):469-73.
- Pesonen AK, Martikainen S, Kajantie E, Heinonen K, Wehkalampi K, Lahti J, Strandberg T, Räikkönen K. The associations between adolescent sleep, diurnal cortisol patterns and cortisol reactivity to dexamethasone suppression test. *Psychoneuroendocrinology*. 2014 Nov;49:150-60.
- Pesonen AK, Sjösten NM, Matthews KA, Heinonen K, Martikainen S, Kajantie E, Tammelin T, Eriksson JG, Strandberg T, Räikkönen K. Temporal associations between daytime physical activity and sleep in children. *PLoS One*. 2011;6(8):e22958.
- Randler C. Differences in sleep and circadian preference between Eastern and Western German adolescents. *Chronobiol Int*. 2008 Jul;25(4):565-75.
- Reid A, Maldonado CC, Baker FC. Sleep behavior of South African adolescents. *Sleep*. 2002 Jun 15;25(4):423-7.
- Rey-López JP, de Carvalho HB, de Moraes AC, Ruiz JR, Sjöström M, Marcos A, Polito A, Gottrand F, Manios Y, Kafatos A, Molnar D, Widhalm K, De Henauw S, Moreno LA; HELENA study. Sleep time and cardiovascular risk factors in adolescents: the HELENA (Healthy Lifestyle in Europe by Nutrition in Adolescence) study. *Sleep Med*. 2014 Jan;15(1):104-10.
- Roberts RE, Duong HT. The prospective association between sleep deprivation and depression among adolescents. *Sleep*. 2014 Feb 1;37(2):239-44.
- Russo PM, Bruni O, Lucidi F, Ferri R, Violani C. Sleep habits and circadian preference in Italian children and adolescents. *J Sleep Res*. 2007 Jun;16(2):163-9.
- Räikkönen K, Matthews KA, Pesonen AK, Pyhälä R, Paavonen EJ, Feldt K, Jones A, Phillips DI, Seckl JR, Heinonen K, Lahti J, Komsu N, Järvenpää AL, Eriksson JG, Strandberg TE, Kajantie E. Poor sleep and altered hypothalamic-pituitary-adrenocortical and sympathetic-adrenal-medullary system activity in children. *J Clin Endocrinol Metab*. 2010 May;95(5):2254-61.
- Sadeh A, Gruber R, Raviv A. The effects of sleep restriction and extension on school-age children: what a difference an hour makes. *Child Dev*. 2003 Mar-Apr;74(2):444-55.
- Sadeh A, Gruber R, Raviv A. Sleep, neurobehavioral functioning, and behavior problems in school-age children. *Child Dev*. 2002 Mar-Apr;73(2):405-17.
- Schreck KA, Mulick JA, Smith AF. Sleep problems as possible predictors of intensified symptoms of autism. *Res Dev Disabil*. 2004 Jan-Feb;25(1):57-66.
- Shan XY, Xi B, Cheng H, Hou DQ, Wang Y, Mi J. Prevalence and behavioral risk factors of overweight and obesity among children aged 2-18 in Beijing, China. *Int J Pediatr Obes*. 2010 Oct;5(5):383-9.
- Shochat T, Cohen-Zion M, Tzischinsky O. Functional consequences of inadequate sleep in adolescents: a systematic review. *Sleep Med Rev*. 2014 Feb;18(1):75-87.
- Shochat T, Flint-Bretler O, Tzischinsky O. Sleep patterns, electronic media exposure and daytime sleep-related behaviours among Israeli adolescents. *Acta Paediatr*. 2010 Sep;99(9):1396-400.
- Short MA, Gradisar M, Lack LC, Wright HR, Dohnt H. The sleep patterns and well-being of Australian adolescents. *J Adolesc*. 2013 Feb;36(1):103-10.
- Short MA, Gradisar M, Lack LC, Wright HR. The impact of sleep on adolescent depressed mood, alertness and academic performance. *J Adolesc*. 2013 Dec;36(6):1025-33.
- Shur-Fen Gau S. Prevalence of sleep problems and their association with inattention/hyperactivity among children aged 6-15 in Taiwan. *J Sleep Res*. 2006 Dec;15(4):403-14. Erratum in: *J Sleep Res*. 2010 Jun;19(2):379. Shur-Fen, Gau Susan [corrected to Shur-Fen Gau, Susan].
- Sivertsen B, Harvey AG, Reichborn-Kjennerud T, Torgersen L, Ystrom E, Hysing M. Later emotional and behavioral problems associated with sleep problems in toddlers: a longitudinal study. *JAMA Pediatr*. 2015 Jun;169(6):575-82.
- Sivertsen B, Harvey AG, Lundervold AJ, Hysing M. Sleep problems and depression in adolescence: results from a large population-based study of Norwegian adolescents aged 16-18 years. *Eur Child Adolesc Psychiatry*. 2014 Aug;23(8):681-9.
- Skidmore PM, Howe AS, Polak MA, Wong JE, Lubransky A, Williams SM, Black KE. Sleep duration and adiposity in older adolescents from Otago, New Zealand: relationships differ between boys and girls and are independent of food choice. *Nutr J*. 2013 Sep 14;12:128.
- Souders MC, Mason TB, Valladares O, Bucan M, Levy SE, Mandell DS, Weaver TE, Pinto-Martin J. Sleep behaviors and sleep quality in children with autism spectrum disorders. *Sleep*. 2009 Dec;32(12):1566-78.
- Spilsbury JC, Storfer-Isser A, Drotar D, Rosen CL, Kirchner HL, Redline S. Effects of the home environment on school-aged children's sleep. *Sleep*. 2005 Nov;28(11):1419-27.
- Spilsbury JC, Storfer-Isser A, Drotar D, Rosen CL, Kirchner LH, Benham H, Redline S. Sleep behavior in an urban US sample of school-aged children. *Arch Pediatr Adolesc Med*. 2004 Oct;158(10):988-94.
- Spruyt K, Aitken RJ, So K, Charlton M, Adamson TM, Horne RS. Relationship between sleep/wake patterns, temperament and overall development in term infants over the first year of life. *Early Hum Dev*. 2008 May;84(5):289-96.
- Stea TH, Knutsen T, Torstveit MK. Association between short time in bed, health-risk behaviors and poor academic achievement among Norwegian adolescents. *Sleep Med*. 2014 Jun;15(6):666-71.
- Steele RM, van Sluijs EM, Cassidy A, Griffin SJ, Ekelund U. Targeting sedentary time or moderate- and vigorous-intensity activity: independent relations with adiposity in a population-based sample of 10-y-old British children. *Am J Clin Nutr*. 2009 Nov;90(5):1185-92.

- Steenari MR, Vuontela V, Paavonen EJ, Carlson S, Fjallberg M, Aronen E. Working memory and sleep in 6- to 13-year-old schoolchildren. *J Am Acad Child Adolesc Psychiatry*. 2003 Jan;42(1):85-92.
- Sugimori H, Yoshida K, Izuno T, Miyakawa M, Suka M, Sekine M, Yamagami T, Kagamimori S. Analysis of factors that influence body mass index from ages 3 to 6 years: A study based on the Toyama cohort study. *Pediatr Int*. 2004 Jun;46(3):302-10.
- Sun Y, Sekine M, Kagamimori S. Lifestyle and overweight among Japanese adolescents: the Toyama Birth Cohort Study. *J Epidemiol*. 2009;19(6):303-10.
- Suzuki H, Kaneita Y, Osaki Y, Minowa M, Kanda H, Suzuki K, Wada K, Hayashi K, Tanihata T, Ohida T. Clarification of the factor structure of the 12-item General Health Questionnaire among Japanese adolescents and associated sleep status. *Psychiatry Res*. 2011 Jun 30;188(1):138-46.
- Takemura T, Funaki K, Kanbayashi T, Kawamoto K, Tsutsui K, Saito Y, Aizawa R, Inomata S, Shimizu T. Sleep habits of students attending elementary schools, and junior and senior high schools in Akita prefecture. *Psychiatry Clin Neurosci*. 2002 Jun;56(3):241-2.
- Tatone-Tokuda F, Dubois L, Ramsay T, Girard M, Touchette E, Petit D, Montplaisir JY. Sex differences in the association between sleep duration, diet and body mass index: a birth cohort study. *J Sleep Res*. 2012 Aug;21(4):448-60.
- Taylor DJ, Jenni OG, Acebo C, Carskadon MA. Sleep tendency during extended wakefulness: insights into adolescent sleep regulation and behavior. *J Sleep Res*. 2005 Sep;14(3):239-44.
- Taylor RW, Williams SM, Farmer VL, Taylor BJ. The stability of sleep patterns in children 3 to 7 years of age. *J Pediatr*. 2015 Mar;166(3):697-702.e1.
- Teixeira LR, Fischer FM, de Andrade MM, Louzada FM, Nagai R. Sleep patterns of day-working, evening high-schooled adolescents of SÃ£o Paulo, Brazil. *Chronobiol Int*. 2004 Mar;21(2):239-52.
- Thorpe K, Staton S, Sawyer E, Pattinson C, Haden C, Smith S. Napping, development and health from 0 to 5 years: a systematic review. *Arch Dis Child*. 2015 Jul;100(7):615-22.
- Touchette E, Côté SM, Petit D, Liu X, Boivin M, Falissard B, Tremblay RE, Montplaisir JY. Short nighttime sleep-duration and hyperactivity trajectories in early childhood. *Pediatrics*. 2009 Nov;124(5):e985-93.
- Unalan D, Ozturk A, Ismailogullari S, Akgul N, Aksu M. The effect of sleep duration and quality on academical success of the elementary school children in Kayseri Turkey. *J Pak Med Assoc*. 2013 May;63(5):576-80.
- van der Vinne V, Zerbini G, Siersema A, Pieper A, Merrow M, Hut RA, Roenneberg T, Kantermann T. Timing of examinations affects school performance differently in early and late chronotypes. *J Biol Rhythms*. 2015 Feb;30(1):53-60.
- Vazsonyi AT, Harris C, Terveer AM, Pagava K, Phagava H, Michaud PA. Parallel mediation effects by sleep on the parental warmth-problem behavior links: evidence from national probability samples of Georgian and Swiss adolescents. *J Youth Adolesc*. 2015 Feb;44(2):331-45.
- Verrillo E, Bizzarri C, Cappa M, Bruni O, Pavone M, Ferri R, Cutrera R. Sleep characteristics in children with growth hormone deficiency. *Neuroendocrinology*. 2011;94(1):66-74.
- Voderholzer U, Piosczyk H, Holz J, Landmann N, Feige B, Loessl B, Kopasz M, Doerr JP, Riemann D, Nissen C. Sleep restriction over several days does not affect long-term recall of declarative and procedural memories in adolescents. *Sleep Med*. 2011 Feb;12(2):170-8.
- Vriend JL, Davidson FD, Corkum PV, Rusak B, Chambers CT, McLaughlin EN. Manipulating sleep duration alters emotional functioning and cognitive performance in children. *J Pediatr Psychol*. 2013 Nov;38(10):1058-69.
- Waldie KE, Thompson JM, Mia Y, Murphy R, Wall C, Mitchell EA. Risk factors for migraine and tension-type headache in 11 year old children. *J Headache Pain*. 2014 Sep 10;15:60.
- Wang H, Sekine M, Chen X, Yamagami T, Kagamimori S. Lifestyle at 3 years of age and quality of life (QOL) in first-year junior high school students in Japan: results of the Toyama Birth Cohort Study. *Qual Life Res*. 2008 Mar;17(2):257-65.
- Weiss A, Xu F, Storfer-Isser A, Thomas A, levers-Landis CE, Redline S. The association of sleep duration with adolescents' fat and carbohydrate consumption. *Sleep*. 2010 Sep;33(9):1201-9.
- Weissbluth M, Davis AT, Poncher J. Night waking in 4- to 8-month-old infants. *J Pediatr*. 1984 Mar;104(3):477-80.
- Weissbluth M, Brouillette RT, Liu K, Hunt CE. Sleep apnea, sleep duration, and infant temperament. *J Pediatr*. 1982 Aug;101(2):307-10.
- Weissbluth M. Sleep duration and infant temperament. *J Pediatr*. 1981 Nov;99(5):817-9.
- Wells JC, Hallal PC, Reichert FF, Menezes AM, Araújo CL, Victora CG. Sleep patterns and television viewing in relation to obesity and blood pressure: evidence from an adolescent Brazilian birth cohort. *Int J Obes (Lond)*. 2008 Jul;32(7):1042-9.
- Westerlund L, Ray C, Roos E. Associations between sleeping habits and food consumption patterns among 10-11-year-old children in Finland. *Br J Nutr*. 2009 Nov;102(10):1531-7.
- Winsler A, Deutsch A, Vorona RD, Payne PA, Szklo-Coxe M. Sleepless in Fairfax: the difference one more hour of sleep can make for teen hopelessness, suicidal ideation, and substance use. *J Youth Adolesc*. 2015 Feb;44(2):362-78.
- Yang CK, Kim JK, Patel SR, Lee JH. Age-related changes in sleep/wake patterns among Korean teenagers. *Pediatrics*. 2005 Jan;115(1 Suppl):250-6.
- Yen CF, King BH, Tang TC. The association between short and long nocturnal sleep durations and risky behaviours and the moderating factors in Taiwanese adolescents. *Psychiatry Res*. 2010 Aug 30;179(1):69-74.
- Yip T. The effects of ethnic/racial discrimination and sleep quality on depressive symptoms and self-esteem trajectories among diverse adolescents. *J Youth Adolesc*. 2015 Feb;44(2):419-30.
- Yokomaku A, Misao K, Omoto F, Yamagishi R, Tanaka K, Takada K, Kohyama J. A study of the association between

sleep habits and problematic behaviors in preschool children. *Chronobiol Int*. 2008 Jul;25(4):549-64.

General Health

Amschler DH, McKenzie JF. Elementary students' sleep habits and teacher observations of sleep-related problems. *J Sch Health*. 2005 Feb;75(2):50-6.

Arman AR, Ay P, Fis NP, Ersu R, Topuzoglu A, Isik U, Berkem M. Association of sleep duration with socio-economic status and behavioural problems among schoolchildren. *Acta Paediatr*. 2011 Mar;100(3):420-4.

Bertelle V, Sevestre A, Laou-Hap K, Nagahapitiye MC, Sizun J. Sleep in the neonatal intensive care unit. *J Perinat Neonatal Nurs*. 2007 Apr-Jun;21(2):140-8; quiz 149-50. Review.

Bertelle V, Mabin D, Adrien J, Sizun J. Sleep of preterm neonates under developmental care or regular environmental conditions. *Early Hum Dev*. 2005 Jul;81(7):595-600.

Byars KC, Yolton K, Rausch J, Lanphear B, Beebe DW. Prevalence, patterns, and persistence of sleep problems in the first 3 years of life. *Pediatrics*. 2012 Feb;129(2):e276-84.

Chugh IM, Khanna P, Shah A. Nocturnal symptoms and sleep disturbances in clinically stable asthmatic children. *Asian Pac J Allergy Immunol*. 2006 Jun-Sep;24(2-3):135-42.

Daniel LC, Boergers J, Kopel SJ, Koinis-Mitchell D. Missed sleep and asthma morbidity in urban children. *Ann Allergy Asthma Immunol*. 2012 Jul;109(1):41-6.

El-Sheikh M, Buckhalt JA, Keller PS, Granger DA. Children's objective and subjective sleep disruptions: links with afternoon cortisol levels. *Health Psychol*. 2008 Jan;27(1):26-33.

Ertan P, Yilmaz O, Caglayan M, Sogut A, Aslan S, Yuksel H. Relationship of sleep quality and quality of life in children with monosymptomatic enuresis. *Child Care Health Dev*. 2009 Jul;35(4):469-74.

Gau SF. Neuroticism and sleep-related problems in adolescence. *Sleep*. 2000 Jun 15;23(4):495-502.

Gerber L. Sleep deprivation in children: a growing public health concern. *Nursing*. 2014 Apr;44(4):50-4. No abstract available.

Hiscock H, Sciberras E, Mensah F, Gerner B, Efron D, Khano S, Oberklaid F. Impact of a behavioural sleep intervention on symptoms and sleep in children with attention deficit hyperactivity disorder, and parental mental health: randomised controlled trial. *BMJ*. 2015 Jan 20;350:h68.

Kostyun RO, Milewski MD, Hafeez I. Sleep disturbance and neurocognitive function during the recovery from a sport-related concussion in adolescents. *Am J Sports Med*. 2015 Mar;43(3):633-40.

Lam LT, Yang L. Short duration of sleep and unintentional injuries among adolescents in China. *Am J Epidemiol*. 2007 Nov 1;166(9):1053-8.

Li Y, Jin H, Owens JA, Hu C. The association between sleep and injury among school-aged children in rural China: a case-control study. *Sleep Med*. 2008 Jan;9(2):142-8.

Lin WH, Yi CC. Unhealthy sleep practices, conduct problems, and daytime functioning during adolescence. *J Youth Adolesc*. 2015 Feb;44(2):431-46.

Magee CA, Lee JK, Vella SA. Bidirectional relationships between sleep duration and screen time in early childhood. *JAMA Pediatr*. 2014 May;168(5):465-70.

Magee CA, Gordon R, Caputi P. Distinct developmental trends in sleep duration during early childhood. *Pediatrics*. 2014 Jun;133(6):e1561-7.

Matricciani L, Blunden S, Rigney G, Williams MT, Olds TS. Children's sleep needs: is there sufficient evidence to recommend optimal sleep for children? *Sleep*. 2013 Apr 1;36(4):527-34.

Matthews KA, Hall M, Dahl RE. Sleep in healthy black and white adolescents. *Pediatrics*. 2014 May;133(5):e1189-96.

McHale SM, Kim JY, Kan M, Updegraff KA. Sleep in Mexican-American adolescents: social ecological and well-being correlates. *J Youth Adolesc*. 2011 Jun;40(6):666-79.

Milewski MD, Skaggs DL, Bishop GA, Pace JL, Ibrahim DA, Wren TA, Barzdukas A. Chronic lack of sleep is associated with increased sports injuries in adolescent athletes. *J Pediatr Orthop*. 2014 Mar;34(2):129-33.

Olds T, Blunden S, Petkov J, Forchino F. The relationships between sex, age, geography and time in bed in adolescents: a meta-analysis of data from 23 countries. *Sleep Med Rev*. 2010 Dec;14(6):371-8.

Owens JA, Fernando S, Mc Guinn M. Sleep disturbance and injury risk in young children. *Behav Sleep Med*. 2005;3(1):18-31.

Paavonen EJ, Nieminen-von Wendt T, Vanhala R, Aronen ET, von Wendt L. Effectiveness of melatonin in the treatment of sleep disturbances in children with Asperger disorder. *J Child Adolesc Psychopharmacol*. 2003 Spring;13(1):83-95.

Park YM, Matsumoto K, Seo YJ, Shinkoda H. Sleep and chronotype for children in Japan. *Percept Mot Skills*. 1999 Jun;88(3 Pt 2):1315-29.

Scott N, Blair PS, Emond AM, Fleming PJ, Humphreys JS, Henderson J, Gringras P. Sleep patterns in children with ADHD: a population-based cohort study from birth to 11 years. *J Sleep Res*. 2013 Apr;22(2):121-8.

Segura-Jiménez V, Carbonell-Baeza A, Keating XD, Ruiz JR, Castro-Piñero J. Association of sleep patterns with psychological positive health and health complaints in children and adolescents. *Qual Life Res*. 2015 Apr;24(4):885-95.

Serra-Negra JM, Paiva SM, Fulgêncio LB, Chavez BA, Lage CF, Pordeus IA. Environmental factors, sleep duration, and sleep bruxism in Brazilian schoolchildren: a case-control study. *Sleep Med*. 2014 Feb;15(2):236-9.

Shi Z, Taylor AW, Gill TK, Tuckerman J, Adams R, Martin J. Short sleep duration and obesity among Australian children. *BMC Public Health*. 2010 Oct 15;10:609.

Short MA, Gradisar M, Lack LC, Wright HR, Dohnt H. The sleep patterns and well-being of Australian adolescents. *J Adolesc*. 2013 Feb;36(1):103-10.

Smaldone A, Honig JC, Byrne MW. Sleepless in America: inadequate sleep and relationships to health and well-

being of our nation's children. *Pediatrics*. 2007 Feb;119 Suppl 1:S29-37.

Spruyt K, Alaribe CU, Nwabara OU. To sleep or not to sleep: a repeated daily challenge for African American children. *CNS Neurosci Ther*. 2015 Jan;21(1):23-31.

Stallones L, Beseler C, Chen P. Sleep patterns and risk of injury among adolescent farm residents. *Am J Prev Med*. 2006 Apr;30(4):300-4.

Valent F, Brusaferrero S, Barbone F. A case-crossover study of sleep and childhood injury. *Pediatrics*. 2001 Feb;107(2):E23.

van der Vinne V, Zerbini G, Siersema A, Pieper A, Merrow M, Hut RA, Roenneberg T, Kantermann T. Timing of examinations affects school performance differently in early and late chronotypes. *J Biol Rhythms*. 2015 Feb;30(1):53-60.

Vazsonyi AT, Harris C, Terveer AM, Pagava K, Phagava H, Michaud PA. Parallel mediation effects by sleep on the parental warmth-problem behavior links: evidence from national probability samples of Georgian and Swiss adolescents. *J Youth Adolesc*. 2015 Feb;44(2):331-45.

Wang H, Sekine M, Chen X, Yamagami T, Kagamimori S. Lifestyle at 3 years of age and quality of life (QOL) in first-year junior high school students in Japan: results of the Toyama Birth Cohort Study. *Qual Life Res*. 2008 Mar;17(2):257-65.

Willoughby MT, Angold A, Egger HL. Parent-reported attention-deficit/hyperactivity disorder symptomatology and sleep problems in a preschool-age pediatric clinic sample. *J Am Acad Child Adolesc Psychiatry*. 2008 Sep;47(9):1086-94.

Human Performance

Abe T, Hagihara A, Nobutomo K. Sleep patterns and impulse control among Japanese junior high school students. *J Adolesc*. 2010 Oct;33(5):633-41.

Amschler DH, McKenzie JF. Elementary students' sleep habits and teacher observations of sleep-related problems. *J Sch Health*. 2005 Feb;75(2):50-6.

Anderson B, Storfer-Isser A, Taylor HG, Rosen CL, Redline S. Associations of executive function with sleepiness and sleep duration in adolescents. *Pediatrics*. 2009 Apr;123(4):e701-7.

Andrade MM, Benedito-Silva AA, Domenice S, Arnhold IJ, Menna-Barreto L. Sleep characteristics of adolescents: a longitudinal study. *J Adolesc Health*. 1993 Jul;14(5):401-6.

Arakawa M, Taira K, Tanaka H, Yamakawa K, Toguchi H, Kadekaru H, Yamamoto Y, Uezu E, Shirakawa S. A survey of junior high school students' sleep habit and lifestyle in Okinawa. *Psychiatry Clin Neurosci*. 2001 Jun;55(3):211-2.

Astill RG, Van der Heijden KB, Van Ijzendoorn MH, Van Someren EJ. Sleep, cognition, and behavioral problems in school-age children: a century of research meta-analyzed. *Psychol Bull*. 2012 Nov;138(6):1109-38.

Beebe DW, Ris MD, Kramer ME, Long E, Amin R. The association between sleep disordered breathing, academic grades, and cognitive and behavioral functioning among overweight subjects during middle to late childhood. *Sleep*. 2010 Nov;33(11):1447-56.

Beebe DW, Rose D, Amin R. Attention, learning, and arousal of experimentally sleep-restricted adolescents in a simulated classroom. *J Adolesc Health*. 2010 Nov;47(5):523-5.

Bell JF, Zimmerman FJ. Shortened nighttime sleep duration in early life and subsequent childhood obesity. *Arch Pediatr Adolesc Med*. 2010 Sep;164(9):840-5. Erratum in: *Arch Pediatr Adolesc Med*. 2010 Nov;164(11):1070.

Blunden SL, Beebe DW. The contribution of intermittent hypoxia, sleep debt and sleep disruption to daytime performance deficits in children: consideration of respiratory and non-respiratory sleep disorders. *Sleep Med Rev*. 2006 Apr;10(2):109-18.

Boto LR, Crispim JN, de Melo IS, Juvandes C, Rodrigues T, Azeredo P, Ferreira R. Sleep deprivation and accidental fall risk in children. *Sleep Med*. 2012 Jan;13(1):88-95.

Buckhalt JA, El-Sheikh M, Keller P. Children's sleep and cognitive functioning: race and socioeconomic status as moderators of effects. *Child Dev*. 2007 Jan-Feb;78(1):213-31.

Carskadon MA. Sleep's effects on cognition and learning in adolescence. *Prog Brain Res*. 2011;190:137-43.

Chung KF, Cheung MM. Sleep-wake patterns and sleep disturbance among Hong Kong Chinese adolescents. *Sleep*. 2008 Feb;31(2):185-94.

Cortese S, Konofal E, Yateman N, Mouren MC, Lecendreux M. Sleep and alertness in children with attention-deficit/hyperactivity disorder: a systematic review of the literature. *Sleep*. 2006 Apr;29(4):504-11. Review.

Countryman AJ, Saab PG, Llabre MM, Penedo FJ, McCalla JR, Schneiderman N. Cardiometabolic risk in adolescents: associations with physical activity, fitness, and sleep. *Ann Behav Med*. 2013 Feb;45(1):121-31.

Dahl RE. Biological, developmental, and neurobehavioral factors relevant to adolescent driving risks. *Am J Prev Med*. 2008 Sep;35(3 Suppl):S278-84.

Dahl RE, Ryan ND, Matty MK, Birmaher B, al-Shabbout M, Williamson DE, Kupfer DJ. Sleep onset abnormalities in depressed adolescents. *Biol Psychiatry*. 1996 Mar 15;39(6):400-10.

Davis AL, Avis KT, Schwebel DC. The effects of acute sleep restriction on adolescents' pedestrian safety in a virtual environment. *J Adolesc Health*. 2013 Dec;53(6):785-90.

El-Sheikh M, Buckhalt JA, Mark Cummings E, Keller P. Sleep disruptions and emotional insecurity are pathways of risk for children. *J Child Psychol Psychiatry*. 2007 Jan;48(1):88-96.

Escribano C, Díaz-Morales JF. Daily fluctuations in attention at school considering starting time and chronotype: an exploratory study. *Chronobiol Int*. 2014 Jul;31(6):761-9.

Fallone G, Acebo C, Seifer R, Carskadon MA. Experimental restriction of sleep opportunity in children: effects on teacher ratings. *Sleep*. 2005 Dec;28(12):1561-7.

Franco P, Seret N, Van Hees JN, Lanquart JP Jr, Groswasser J, Kahn A. Cardiac changes during sleep in sleep-deprived infants. *Sleep*. 2003 Nov 1;26(7):845-8.

Fukuda K, Sakashita Y. Sleeping pattern of kindergartners and nursery school children: function of daytime nap. *Percept Mot Skills*. 2002 Feb;94(1):219-28.

- Garmy P, Nyberg P, Jakobsson U. Sleep and television and computer habits of Swedish school-age children. *J Sch Nurs*. 2012 Dec;28(6):469-76.
- Gau SF. Neuroticism and sleep-related problems in adolescence. *Sleep*. 2000 Jun 15;23(4):495-502.
- Gau SS, Shang CY, Merikangas KR, Chiu YN, Soong WT, Cheng AT. Association between morningness-eveningness and behavioral/emotional problems among adolescents. *J Biol Rhythms*. 2007 Jun;22(3):268-74.
- Geiger A, Achermann P, Jenni OG. Association between sleep duration and intelligence scores in healthy children. *Dev Psychol*. 2010 Jul;46(4):949-54.
- Gibbs S, Wiltshire E, Elder D. Nocturnal sleep measured by actigraphy in children with Prader-Willi syndrome. *J Pediatr*. 2013 Apr;162(4):765-9.
- Gibson ES, Powles AC, Thabane L, O'Brien S, Molnar DS, Trajanovic N, Ogilvie R, Shapiro C, Yan M, Chilcott-Tanser L. "Sleepiness" is serious in adolescence: two surveys of 3235 Canadian students." *BMC Public Health*. 2006 May 2;6:116.
- Greenfeld M, Constantini S, Tauman R, Sivan Y. Sleep disturbances in children recovered from central nervous system neoplasms. *J Pediatr*. 2011 Aug;159(2):268-72.e1.
- Gruber R, Cassoff J, Frenette S, Wiebe S, Carrier J. Impact of sleep extension and restriction on children's emotional lability and impulsivity. *Pediatrics*. 2012 Nov;130(5):e1155-61.
- Gruber R, Wiebe S, Montecalvo L, Brunetti B, Amsel R, Carrier J. Impact of sleep restriction on neurobehavioral functioning of children with attention deficit hyperactivity disorder. *Sleep*. 2011 Mar 1;34(3):315-23.
- Gruber R, Laviolette R, Deluca P, Monson E, Cornish K, Carrier J. Short sleep duration is associated with poor performance on IQ measures in healthy school-age children. *Sleep Med*. 2010 Mar;11(3):289-94.
- Guérin N, Reinberg A, Testu F, Boulenguez S, Mechkouri M, Touitou Y. Role of school schedule, age, and parental socioeconomic status on sleep duration and sleepiness of Parisian children. *Chronobiol Int*. 2001 Nov;18(6):1005-17.
- Hansen M, Janssen I, Schiff A, Zee PC, Dubocovich ML. The impact of school daily schedule on adolescent sleep. *Pediatrics*. 2005 Jun;115(6):1555-61.
- Henderson JM, France KG, Owens JL, Blampied NM. Sleeping through the night: the consolidation of self-regulated sleep across the first year of life. *Pediatrics*. 2010 Nov;126(5):e1081-7.
- Hjorth MF, Chaput JP, Ritz C, Dalskov SM, Andersen R, Astrup A, Tetens I, Michaelsen KF, Sjodin A. Fatness predicts decreased physical activity and increased sedentary time, but not vice versa: support from a longitudinal study in 8- to 11-year-old children. *Int J Obes (Lond)*. 2014 Jul;38(7):959-65.
- Holm SM, Forbes EE, Ryan ND, Phillips ML, Tarr JA, Dahl RE. Reward-related brain function and sleep in pre/early pubertal and mid/late pubertal adolescents. *J Adolesc Health*. 2009 Oct;45(4):326-34.
- Ishihara K. The effect of 2-h sleep reduction by a delayed bedtime on daytime sleepiness in children. *Psychiatry Clin Neurosci*. 1999 Apr;53(2):113-5.
- Ishihara K, Miyake S. A longitudinal study of the development of daytime sleepiness in children. *Psychiatry Clin Neurosci*. 1998 Apr;52(2):178-81.
- Keller PS, El-Sheikh M, Buckhalt JA. Children's attachment to parents and their academic functioning: sleep disruptions as moderators of effects. *J Dev Behav Pediatr*. 2008 Dec;29(6):441-9.
- Kopasz M, Loessl B, Valerius G, Koenig E, Matthaes N, Hornyak M, Kloepfer C, Nissen C, Riemann D, Voderholzer U. No persisting effect of partial sleep curtailment on cognitive performance and declarative memory recall in adolescents. *J Sleep Res*. 2010 Mar;19(1 Pt 1):71-9.
- Kostyun RO, Milewski MD, Hafeez I. Sleep disturbance and neurocognitive function during the recovery from a sport-related concussion in adolescents. *Am J Sports Med*. 2015 Mar;43(3):633-40.
- Lam JC, Mahone EM, Mason T, Scharf SM. The effects of napping on cognitive function in preschoolers. *J Dev Behav Pediatr*. 2011 Feb-Mar;32(2):90-7.
- Lee JA, Park HS. Relation between sleep duration, overweight, and metabolic syndrome in Korean adolescents. *Nutr Metab Cardiovasc Dis*. 2014 Jan;24(1):65-71.
- Leprout R, Van Cauter E. Role of sleep and sleep loss in hormonal release and metabolism. *Endocr Dev*. 2010;17:11-21.
- Lewandowski AS, Palermo TM, De la Motte S, Fu R. Temporal daily associations between pain and sleep in adolescents with chronic pain versus healthy adolescents. *Pain*. 2010 Oct;151(1):220-5.
- Li S, Arguelles L, Jiang F, Chen W, Jin X, Yan C, Tian Y, Hong X, Qian C, Zhang J, Wang X, Shen X. Sleep, school performance, and a school-based intervention among school-aged children: a sleep series study in China. *PLoS One*. 2013 Jul 10;8(7):e67928.
- Lin WH, Yi CC. Unhealthy sleep practices, conduct problems, and daytime functioning during adolescence. *J Youth Adolesc*. 2015 Feb;44(2):431-46.
- Liu X. Sleep and adolescent suicidal behavior. *Sleep*. 2004 Nov 1;27(7):1351-8.
- Mak KK, Lee SL, Ho SY, Lo WS, Lam TH. Sleep and academic performance in Hong Kong adolescents. *J Sch Health*. 2012 Nov;82(11):522-7.
- Martikainen S, Pesonen AK, Feldt K, Jones A, Lahti J, Pyhälä R, Heinonen K, Kajantie E, Eriksson J, Räikkönen K. Poor sleep and cardiovascular function in children. *Hypertension*. 2011 Jul;58(1):16-21.
- Martinez-Gomez D, Eisenmann JC, Gomez-Martinez S, Hill EE, Zapatera B, Veiga OL, Marcos A; AFINOS Study Group. Sleep duration and emerging cardiometabolic risk markers in adolescents. The AFINOS study. *Sleep Med*. 2011 Dec;12(10):997-1002.
- Maski KP, Kothare SV. Sleep deprivation and neurobehavioral functioning in children. *Int J Psychophysiol*. 2013 Aug;89(2):259-64.
- Matsushita M, Koyama A, Ushijima H, Mikami A, Katsumata Y, Kikuchi Y, Ichimi N, Jono T, Fujise N, Ikeda M. Sleep duration and its association with sleepiness and

- depression in ronin-sei" preparatory school students." *Asian J Psychiatr.* 2014 Jun;9:61-6.
- McLaughlin Crabtree V, Beal Korhonen J, Montgomery-Downs HE, Faye Jones V, O'Brien LM, Gozal D. Cultural influences on the bedtime behaviors of young children. *Sleep Med.* 2005 Jul;6(4):319-24.
- Meijer AM. Chronic sleep reduction, functioning at school and school achievement in preadolescents. *J Sleep Res.* 2008 Dec;17(4):395-405. Erratum in: *J Sleep Res.* 2009 Mar;18(1):144.
- Mercer PW, Merritt SL, Cowell JM. Differences in reported sleep need among adolescents. *J Adolesc Health.* 1998 Nov;23(5):259-63.
- Mercier L, Pivik RT, Busby K. Sleep patterns in reading disabled children. *Sleep.* 1993 Apr;16(3):207-15. Review.
- Michels N, Clays E, De Buyzere M, Vanaelst B, De Henauw S, Sioen I. Children's sleep and autonomic function: low sleep quality has an impact on heart rate variability. *Sleep.* 2013 Dec 1;36(12):1939-46.
- Molfese DL, Ivanenko A, Key AF, Roman A, Molfese VJ, O'Brien LM, Gozal D, Kota S, Hudac CM. A one-hour sleep restriction impacts brain processing in young children across tasks: evidence from event-related potentials. *Dev Neuropsychol.* 2013;38(5):317-36.
- Ng EP, Ng DK, Chan CH. Sleep duration, wake/sleep symptoms, and academic performance in Hong Kong Secondary School Children. *Sleep Breath.* 2009 Nov;13(4):357-67.
- Nugent R, Althouse A, Yaqub Y, Nugent K, Raj R. Modeling the relation between obesity and sleep parameters in children referred for dietary weight reduction intervention. *South Med J.* 2014 Aug;107(8):473-80.
- Olds T, Blunden S, Petkov J, Forchino F. The relationships between sex, age, geography and time in bed in adolescents: a meta-analysis of data from 23 countries. *Sleep Med Rev.* 2010 Dec;14(6):371-8.
- Ortega FB, Chillón P, Ruiz JR, Delgado M, Albers U, Alvarez-Granda JL, Marcos A, Moreno LA, Castillo MJ. Sleep patterns in Spanish adolescents: associations with TV watching and leisure-time physical activity. *Eur J Appl Physiol.* 2010 Oct;110(3):563-73.
- Ortega FB, Ruiz JR, Castillo R, Chillón P, Labayen I, Martínez-Gómez D, Redondo C, Marcos A, Moreno LA; AVENA study group. Sleep duration and cognitive performance in adolescence. The AVENA study. *Acta Paediatr.* 2010 Mar;99(3):454-6.
- Owens JA, Belon K, Moss P. Impact of delaying school start time on adolescent sleep, mood, and behavior. *Arch Pediatr Adolesc Med.* 2010 Jul;164(7):608-14.
- Owens JA, Fernando S, Mc Guinn M. Sleep disturbance and injury risk in young children. *Behav Sleep Med.* 2005;3(1):18-31.
- Perez-Lloret S, Videla AJ, Richaudeau A, Vigo D, Rossi M, Cardinali DP, Perez-Chada D. A multi-step pathway connecting short sleep duration to daytime somnolence, reduced attention, and poor academic performance: an exploratory cross-sectional study in teenagers. *J Clin Sleep Med.* 2013 May 15;9(5):469-73.
- Perkinson-Gloor N, Lemola S, Grob A. Sleep duration, positive attitude toward life, and academic achievement: the role of daytime tiredness, behavioral persistence, and school start times. *J Adolesc.* 2013 Apr;36(2):311-8.
- Pesonen AK, Martikainen S, Kajantie E, Heinonen K, Wehkalampi K, Lahti J, Strandberg T, Räikkönen K. The associations between adolescent sleep, diurnal cortisol patterns and cortisol reactivity to dexamethasone suppression test. *Psychoneuroendocrinology.* 2014 Nov;49:150-60.
- Pizza F, Contardi S, Antognini AB, Zagoraiou M, Borrotti M, Mostacci B, Mondini S, Cirignotta F. Sleep quality and motor vehicle crashes in adolescents. *J Clin Sleep Med.* 2010 Feb 15;6(1):41-5.
- Reid A, Maldonado CC, Baker FC. Sleep behavior of South African adolescents. *Sleep.* 2002 Jun 15;25(4):423-7.
- Rey-López JP, de Carvalho HB, de Moraes AC, Ruiz JR, Sjöström M, Marcos A, Polito A, Gottrand F, Manios Y, Kafatos A, Molnar D, Widhalm K, De Henauw S, Moreno LA; HELENA study. Sleep time and cardiovascular risk factors in adolescents: the HELENA (Healthy Lifestyle in Europe by Nutrition in Adolescence) study. *Sleep Med.* 2014 Jan;15(1):104-10.
- Roberts RE, Roberts CR, Duong HT. Sleepless in adolescence: prospective data on sleep deprivation, health and functioning. *J Adolesc.* 2009 Oct;32(5):1045-57.
- Sadeh A, Gruber R, Raviv A. The effects of sleep restriction and extension on school-age children: what a difference an hour makes. *Child Dev.* 2003 Mar-Apr;74(2):444-55.
- Sadeh A, Gruber R, Raviv A. Sleep, neurobehavioral functioning, and behavior problems in school-age children. *Child Dev.* 2002 Mar-Apr;73(2):405-17.
- Sallinen BJ, Hassan F, Olszewski A, Maupin A, Hoban TF, Chervin RD, Woolford SJ. Longer weekly sleep duration predicts greater 3-month BMI reduction among obese adolescents attending a clinical multidisciplinary weight management program. *Obes Facts.* 2013;6(3):239-46.
- Salzarulo P, Fagioli I. Sleep for development or development for waking?--some speculations from a human perspective. *Behav Brain Res.* 1995 Jul-Aug;69(1-2):23-7. Review.
- Sampei M, Murata K, Dakeishi M, Wood DC. Cardiac autonomic hypofunction in preschool children with short nocturnal sleep. *Tohoku J Exp Med.* 2006 Mar;208(3):235-42.
- Segura-Jiménez V, Carbonell-Baeza A, Keating XD, Ruiz JR, Castro-Piñero J. Association of sleep patterns with psychological positive health and health complaints in children and adolescents. *Qual Life Res.* 2015 Apr;24(4):885-95.
- Shochat T, Flint-Bretler O, Tzischinsky O. Sleep patterns, electronic media exposure and daytime sleep-related behaviours among Israeli adolescents. *Acta Paediatr.* 2010 Sep;99(9):1396-400.
- Short MA, Gradisar M, Lack LC, Wright HR. The impact of sleep on adolescent depressed mood, alertness and academic performance. *J Adolesc.* 2013 Dec;36(6):1025-33.
- Spicuzza L, Sciuto C, Leonardi S, La Rosa M. Early occurrence of obstructive sleep apnea in infants and

children with cystic fibrosis. *Arch Pediatr Adolesc Med*. 2012 Dec;166(12):1165-9.

Stallones L, Beseler C, Chen P. Sleep patterns and risk of injury among adolescent farm residents. *Am J Prev Med*. 2006 Apr;30(4):300-4.

Steele RM, van Sluijs EM, Cassidy A, Griffin SJ, Ekelund U. Targeting sedentary time or moderate- and vigorous-intensity activity: independent relations with adiposity in a population-based sample of 10-y-old British children. *Am J Clin Nutr*. 2009 Nov;90(5):1185-92.

Steenari MR, Vuontela V, Paavonen EJ, Carlson S, Fjallberg M, Aronen E. Working memory and sleep in 6- to 13-year-old schoolchildren. *J Am Acad Child Adolesc Psychiatry*. 2003 Jan;42(1):85-92.

Taki Y, Hashizume H, Thyreau B, Sassa Y, Takeuchi H, Wu K, Kotozaki Y, Nouchi R, Asano M, Asano K, Fukuda H, Kawashima R. Sleep duration during weekdays affects hippocampal gray matter volume in healthy children. *Neuroimage*. 2012 Mar;60(1):471-5.

Teixeira LR, Fischer FM, Nagai R, Turte SL. Teen at work: the burden of a double shift on daily activities. *Chronobiol Int*. 2004;21(6):845-58.

Touchette E, Petit D, Séguin JR, Boivin M, Tremblay RE, Montplaisir JY. Associations between sleep duration patterns and behavioral/cognitive functioning at school entry. *Sleep*. 2007 Sep;30(9):1213-9.

Unalan D, Ozturk A, Ismailogullari S, Akgul N, Aksu M. The effect of sleep duration and quality on academical success of the elementary school children in Kayseri Turkey. *J Pak Med Assoc*. 2013 May;63(5):576-80.

Urschitz MS, Heine K, Brockmann PE, Peters T, Durst W, Poets CF, Wilhelm B. Subjective and objective daytime sleepiness in schoolchildren and adolescents: results of a community-based study. *Sleep Med*. 2013 Oct;14(10):1005-12.

Valent F, Brusaferrero S, Barbone F. A case-crossover study of sleep and childhood injury. *Pediatrics*. 2001 Feb;107(2):E23.

Verrillo E, Bizzarri C, Cappa M, Bruni O, Pavone M, Ferri R, Cutrera R. Sleep characteristics in children with growth hormone deficiency. *Neuroendocrinology*. 2011;94(1):66-74.

Voderholzer U, Piosczyk H, Holz J, Landmann N, Feige B, Loessel B, Kopasz M, Doerr JP, Riemann D, Nissen C. Sleep restriction over several days does not affect long-term recall of declarative and procedural memories in adolescents. *Sleep Med*. 2011 Feb;12(2):170-8.

Vriend JL, Davidson FD, Corkum PV, Rusak B, Chambers CT, McLaughlin EN. Manipulating sleep duration alters emotional functioning and cognitive performance in children. *J Pediatr Psychol*. 2013 Nov;38(10):1058-69.

Westerlund L, Ray C, Roos E. Associations between sleeping habits and food consumption patterns among 10-11-year-old children in Finland. *Br J Nutr*. 2009 Nov;102(10):1531-7.

Zhu Y, Li AM, Au CT, Kong AP, Zhang J, Wong CK, Chan JC, Wing YK. Association between sleep architecture and glucose tolerance in children and adolescents. *J Diabetes*. 2015 Jan;7(1):10-5.

Immunologic Health

Berentzen NE, Smit HA, Bekkers MB, Brunekreef B, Koppelman GH, De Jongste JC, Kerkhof M, Van Rossem L, Wijga AH. Time in bed, sleep quality and associations with cardiometabolic markers in children: the Prevention and Incidence of Asthma and Mite Allergy birth cohort study. *J Sleep Res*. 2014 Feb;23(1):3-12.

Börnhorst C, Hense S, Ahrens W, Hebestreit A, Reisch L, Barba G, von Kries R, Bayer O; IDEFICS Consortium. From sleep duration to childhood obesity--what are the pathways? *Eur J Pediatr*. 2012 Jul;171(7):1029-38.

Chugh IM, Khanna P, Shah A. Nocturnal symptoms and sleep disturbances in clinically stable asthmatic children. *Asian Pac J Allergy Immunol*. 2006 Jun-Sep;24(2-3):135-42.

Countryman AJ, Saab PG, Llabre MM, Penedo FJ, McCalla JR, Schneiderman N. Cardiometabolic risk in adolescents: associations with physical activity, fitness, and sleep. *Ann Behav Med*. 2013 Feb;45(1):121-31.

Daniel LC, Boergers J, Kopel SJ, Koinis-Mitchell D. Missed sleep and asthma morbidity in urban children. *Ann Allergy Asthma Immunol*. 2012 Jul;109(1):41-6.

El-Sheikh M, Buckhalt JA, Granger DA, Erath SA, Acebo C. The association between children's sleep disruption and salivary interleukin-6. *J Sleep Res*. 2007 Jun;16(2):188-97.

Kieckhefer GM, Ward TM, Tsai SY, Lentz MJ. Nighttime sleep and daytime nap patterns in school age children with and without asthma. *J Dev Behav Pediatr*. 2008 Oct;29(5):338-44.

Martinez-Gomez D, Eisenmann JC, Gomez-Martinez S, Hill EE, Zapatera B, Veiga OL, Marcos A; AFINOS Study Group. Sleep duration and emerging cardiometabolic risk markers in adolescents. The AFINOS study. *Sleep Med*. 2011 Dec;12(10):997-1002.

Moore M, Kirchner HL, Drotar D, Johnson N, Rosen C, Ancoli-Israel S, Redline S. Relationships among sleepiness, sleep time, and psychological functioning in adolescents. *J Pediatr Psychol*. 2009 Nov-Dec;34(10):1175-83.

Moore M, Kirchner HL, Drotar D, Johnson N, Rosen C, Redline S. Correlates of adolescent sleep time and variability in sleep time: the role of individual and health related characteristics. *Sleep Med*. 2011 Mar;12(3):239-45.

Pérez de Heredia F, Garaulet M, Gómez-Martínez S, Díaz LE, Wärnberg J, Androustos O, Michels N, Breidenassel C, Cuenca-García M, Huybrechts I, Gottrand F, Ferrari M, Santaliestra-Pasías AM, Kafatos A, Molnár D, Sjöström M, Widhalm K, Moreno LA, Marcos A; HELENA Study Group. Self-reported sleep duration, white blood cell counts and cytokine profiles in European adolescents: the HELENA study. *Sleep Med*. 2014 Oct;15(10):1251-8.

Spruyt K, Molfese DL, Gozal D. Sleep duration, sleep regularity, body weight, and metabolic homeostasis in school-aged children. *Pediatrics*. 2011 Feb;127(2):e345-52.

Tirosh E, Scher A, Sadeh A, Jaffe M, Lavie P. Sleep characteristics of asthmatics in the first four years of life: a comparative study. *Arch Dis Child*. 1993 Apr;68(4):481-3.

Longevity

- Aishworiya R, Chan P, Kiing J, Chong SC, Laino AG, Tay SKh. Sleep behaviour in a sample of preschool children in Singapore. *Ann Acad Med Singapore*. 2012 Mar;41(3):99-104.
- Azadbakht L, Kelishadi R, Khodarahmi M, Qorbani M, Heshmat R, Motlagh ME, Taslimi M, Ardalan G. The association of sleep duration and cardiometabolic risk factors in a national sample of children and adolescents: the CASPIAN III study. *Nutrition*. 2013 Sep;29(9):1133-41.
- Barnes JC, Meldrum RC. The impact of sleep duration on adolescent development: a genetically informed analysis of identical twin pairs. *J Youth Adolesc*. 2015 Feb;44(2):489-506.
- Beebe DW, Lewin D, Zeller M, McCabe M, MacLeod K, Daniels SR, Amin R. Sleep in overweight adolescents: shorter sleep, poorer sleep quality, sleepiness, and sleep-disordered breathing. *J Pediatr Psychol*. 2007 Jan-Feb;32(1):69-79.
- Berentzen NE, Smit HA, Bekkers MB, Brunekreef B, Koppelman GH, De Jongste JC, Kerkhof M, Van Rossem L, Wijga AH. Time in bed, sleep quality and associations with cardiometabolic markers in children: the Prevention and Incidence of Asthma and Mite Allergy birth cohort study. *J Sleep Res*. 2014 Feb;23(1):3-12.
- Bibiloni Mdel M, Martinez E, Llull R, Juarez MD, Pons A, Tur JA. Prevalence and risk factors for obesity in Balearic Islands adolescents. *Br J Nutr*. 2010 Jan;103(1):99-106.
- Bonuck K, Chervin RD, Howe LD. Sleep-disordered breathing, sleep duration, and childhood overweight: a longitudinal cohort study. *J Pediatr*. 2015 Mar;166(3):632-9.
- Byars KC, Yolton K, Rausch J, Lanphear B, Beebe DW. Prevalence, patterns, and persistence of sleep problems in the first 3 years of life. *Pediatrics*. 2012 Feb;129(2):e276-84.
- Chung KF, Cheung MM. Sleep-wake patterns and sleep disturbance among Hong Kong Chinese adolescents. *Sleep*. 2008 Feb;31(2):185-94.
- Daniel LC, Boergers J, Kopel SJ, Koinis-Mitchell D. Missed sleep and asthma morbidity in urban children. *Ann Allergy Asthma Immunol*. 2012 Jul;109(1):41-6.
- Danielsen YS, Pallesen S, Stormark KM, Nordhus IH, Bjorvatn B. The relationship between school day sleep duration and body mass index in Norwegian children (aged 10-12). *Int J Pediatr Obes*. 2010 May 3;5(3):214-20.
- de Jong E, Stocks T, Visscher TL, HiraSing RA, Seidell JC, Renders CM. Association between sleep duration and overweight: the importance of parenting. *Int J Obes (Lond)*. 2012 Oct;36(10):1278-84.
- Do YK, Shin E, Bautista MA, Foo K. The associations between self-reported sleep duration and adolescent health outcomes: what is the role of time spent on Internet use? *Sleep Med*. 2013 Feb;14(2):195-200.
- Drescher AA, Goodwin JL, Silva GE, Quan SF. Caffeine and screen time in adolescence: associations with short sleep and obesity. *J Clin Sleep Med*. 2011 Aug 15;7(4):337-42.
- Eaton DK, McKnight-Eily LR, Lowry R, Perry GS, Presley-Cantrell L, Croft JB. Prevalence of insufficient, borderline, and optimal hours of sleep among high school students - United States, 2007. *J Adolesc Health*. 2010 Apr;46(4):399-401.
- Elsenburg LK, Corpeleijn E, van Sluijs EM, Atkin AJ. Clustering and correlates of multiple health behaviours in 9-10 year old children. *PLoS One*. 2014;9(6):e99498.
- Fitzgerald CT, Messias E, Buysse DJ. Teen sleep and suicidality: results from the youth risk behavior surveys of 2007 and 2009. *J Clin Sleep Med*. 2011 Aug 15;7(4):351-6.
- Franco P, Seret N, Van Hees JN, Lanquart JP Jr, Groswasser J, Kahn A. Cardiac changes during sleep in sleep-deprived infants. *Sleep*. 2003 Nov 1;26(7):845-8.
- Gau SS, Shang CY, Merikangas KR, Chiu YN, Soong WT, Cheng AT. Association between morningness-eveningness and behavioral/emotional problems among adolescents. *J Biol Rhythms*. 2007 Jun;22(3):268-74.
- Guo X, Zheng L, Li Y, Yu S, Liu S, Zhou X, Zhang X, Sun Z, Wang R, Sun Y. Association between sleep duration and hypertension among Chinese children and adolescents. *Clin Cardiol*. 2011 Dec;34(12):774-81.
- Guo X, Zheng L, Li Y, Yu S, Sun G, Yang H, Zhou X, Zhang X, Sun Z, Sun Y. Differences in lifestyle behaviors, dietary habits, and familial factors among normal-weight, overweight, and obese Chinese children and adolescents. *Int J Behav Nutr Phys Act*. 2012 Oct 2;9:120.
- Gupta R, Bhatia MS, Chhabra V, Sharma S, Dahiya D, Semalti K, Sapra R, Dua RS. Sleep patterns of urban school-going adolescents. *Indian Pediatr*. 2008 Mar;45(3):183-9.
- Hart CN, Cairns A, Jelalian E. Sleep and obesity in children and adolescents. *Pediatr Clin North Am*. 2011 Jun;58(3):715-33.
- Hjorth MF, Quist JS, Andersen R, Michaelsen KF, Tetens I, Astrup A, Chaput JP, Sjödén A. Change in sleep duration and proposed dietary risk factors for obesity in Danish school children. *Pediatr Obes*. 2014 Dec;9(6):e156-9.
- Jaffa T, Scott S, Hendriks JH, Shapiro CM. ABC of sleep disorders. *Sleep disorders in children*. *BMJ*. 1993 Mar 6;306(6878):640-3.
- Jang SI, Lee KS, Park EC. Relationship between current sleep duration and past suicidal ideation or attempt among Korean adolescents. *J Prev Med Public Health*. 2013 Nov;46(6):329-35.
- Kang SG, Lee YJ, Kim SJ, Lim W, Lee HJ, Park YM, Cho IH, Cho SJ, Hong JP. Weekend catch-up sleep is independently associated with suicide attempts and self-injury in Korean adolescents. *Compr Psychiatry*. 2014 Feb;55(2):319-25.
- Kjeldsen JS, Hjorth MF, Andersen R, Michaelsen KF, Tetens I, Astrup A, Chaput JP, Sjödén A. Short sleep duration and large variability in sleep duration are independently associated with dietary risk factors for obesity in Danish school children. *Int J Obes (Lond)*. 2014 Jan;38(1):32-9.
- Kleiser C, Schaffrath Rosario A, Mensink GB, Prinz-Langenohl R, Kurth BM. Potential determinants of obesity among children and adolescents in Germany: results from the cross-sectional KiGGS Study. *BMC Public Health*. 2009 Feb 2;9:46.

- Koren D, O'Sullivan KL, Mokhlesi B. Metabolic and glycemic sequelae of sleep disturbances in children and adults. *Curr Diab Rep.* 2015 Jan;15(1):562.
- Kuciene R, Dulskiene V. Associations of short sleep duration with prehypertension and hypertension among Lithuanian children and adolescents: a cross-sectional study. *BMC Public Health.* 2014 Mar 15;14:255.
- Kwon JA, Lee M, Yoo KB, Park EC. Does the duration and time of sleep increase the risk of allergic rhinitis? Results of the 6-year nationwide Korea youth risk behavior web-based survey. *PLoS One.* 2013;8(8):e72507.
- Lam LT, Yang L. Short duration of sleep and unintentional injuries among adolescents in China. *Am J Epidemiol.* 2007 Nov 1;166(9):1053-8. Epub 2007 Aug 13.
- Lazaratou H, Dikeos DG, Anagnostopoulos DC, Sbokou O, Soldatos CR. Sleep problems in adolescence. A study of senior high school students in Greece. *Eur Child Adolesc Psychiatry.* 2005 Jul;14(4):237-43.
- Lee JA, Park HS. Relation between sleep duration, overweight, and metabolic syndrome in Korean adolescents. *Nutr Metab Cardiovasc Dis.* 2014 Jan;24(1):65-71.
- Lee YJ, Cho SJ, Cho IH, Kim SJ. Insufficient sleep and suicidality in adolescents. *Sleep.* 2012 Apr 1;35(4):455-60.
- Li Y, Jin H, Owens JA, Hu C. The association between sleep and injury among school-aged children in rural China: a case-control study. *Sleep Med.* 2008 Jan;9(2):142-8.
- Liu X, Uchiyama M, Okawa M, Kurita H. Prevalence and correlates of self-reported sleep problems among Chinese adolescents. *Sleep.* 2000 Feb 1;23(1):27-34.
- Liu X. Sleep and adolescent suicidal behavior. *Sleep.* 2004 Nov 1;27(7):1351-8.
- Mak KK, Lee SL, Ho SY, Lo WS, Lam TH. Sleep and academic performance in Hong Kong adolescents. *J Sch Health.* 2012 Nov;82(11):522-7.
- Martikainen S, Pesonen AK, Feldt K, Jones A, Lahti J, Pyhälä R, Heinonen K, Kajantie E, Eriksson J, Räikkönen K. Poor sleep and cardiovascular function in children. *Hypertension.* 2011 Jul;58(1):16-21.
- Matamura M, Tochigi M, Usami S, Yonehara H, Fukushima M, Nishida A, Togo F, Sasaki T. Associations between sleep habits and mental health status and suicidality in a longitudinal survey of monozygotic twin adolescents. *J Sleep Res.* 2014 Jun;23(3):290-4.
- Matricciani LA, Olds TS, Blunden S, Rigney G, Williams MT. Never enough sleep: a brief history of sleep recommendations for children. *Pediatrics.* 2012 Mar;129(3):548-56.
- Mitchell JA, Rodriguez D, Schmitz KH, Audrain-McGovern J. Sleep duration and adolescent obesity. *Pediatrics.* 2013 May;131(5):e1428-34.
- Montgomery-Downs HE, Gozal D. Sleep habits and risk factors for sleep-disordered breathing in infants and young toddlers in Louisville, Kentucky. *Sleep Med.* 2006 Apr;7(3):211-9.
- Owens JA, Maxim R, Nobile C, McGuinn M, Msall M. Parental and self-report of sleep in children with attention-deficit/hyperactivity disorder. *Arch Pediatr Adolesc Med.* 2000 Jun;154(6):549-55.
- Ozturk A, Mazicioglu M, Poyrazoglu S, Cicek B, Gunay O, Kurtoglu S. The relationship between sleep duration and obesity in Turkish children and adolescents. *Acta Paediatr.* 2009 Apr;98(4):699-702.
- Paavonen EJ, Vehkalahti K, Vanhala R, von Wendt L, Nieminen-von Wendt T, Aronen ET. Sleep in children with Asperger syndrome. *J Autism Dev Disord.* 2008 Jan;38(1):41-51.
- Padez C, Mourao I, Moreira P, Rosado V. Long sleep duration and childhood overweight/obesity and body fat. *Am J Hum Biol.* 2009 May-Jun;21(3):371-6.
- Park JH, Yoo JH, Kim SH. Associations between non-restorative sleep, short sleep duration and suicidality: findings from a representative sample of Korean adolescents. *Psychiatry Clin Neurosci.* 2013 Jan;67(1):28-34.
- Pizza F, Contardi S, Antognini AB, Zagoraiou M, Borrotti M, Mostacci B, Mondini S, Cirignotta F. Sleep quality and motor vehicle crashes in adolescents. *J Clin Sleep Med.* 2010 Feb 15;6(1):41-5.
- Rey-López JP, de Carvalho HB, de Moraes AC, Ruiz JR, Sjöström M, Marcos A, Polito A, Gottrand F, Manios Y, Kafatos A, Molnar D, Widhalm K, De Henauw S, Moreno LA; HELENA study. Sleep time and cardiovascular risk factors in adolescents: the HELENA (Healthy Lifestyle in Europe by Nutrition in Adolescence) study. *Sleep Med.* 2014 Jan;15(1):104-10.
- Rutters F, Gerver WJ, Nieuwenhuizen AG, Verhoef SP, Westerterp-Plantenga MS. Sleep duration and body-weight development during puberty in a Dutch children cohort. *Int J Obes (Lond).* 2010 Oct;34(10):1508-14.
- Seicean A, Redline S, Seicean S, Kirchner HL, Gao Y, Sekine M, Zhu X, Storer-Isser A. Association between short sleeping hours and overweight in adolescents: results from a US Suburban High School survey. *Sleep Breath.* 2007 Dec;11(4):285-93.
- Shan XY, Xi B, Cheng H, Hou DQ, Wang Y, Mi J. Prevalence and behavioral risk factors of overweight and obesity among children aged 2-18 in Beijing, China. *Int J Pediatr Obes.* 2010 Oct;5(5):383-9.
- Sivertsen B, Harvey AG, Reichborn-Kjennerud T, Torgersen L, Ystrom E, Hysing M. Later emotional and behavioral problems associated with sleep problems in toddlers: a longitudinal study. *JAMA Pediatr.* 2015 Jun;169(6):575-82.
- Sivertsen B, Pallesen S, Stormark KM, BÅe T, Lundervold AJ, Hysing M. Delayed sleep phase syndrome in adolescents: prevalence and correlates in a large population based study. *BMC Public Health.* 2013 Dec 11;13:1163.
- Skidmore PM, Howe AS, Polak MA, Wong JE, Lubransky A, Williams SM, Black KE. Sleep duration and adiposity in older adolescents from Otago, New Zealand: relationships differ between boys and girls and are independent of food choice. *Nutr J.* 2013 Sep 14;12:128.
- Smaldone A, Honig JC, Byrne MW. Sleepless in America: inadequate sleep and relationships to health and well-being of our nation's children. *Pediatrics.* 2007 Feb;119 Suppl 1:S29-37.
- Souders MC, Mason TB, Valladares O, Bucan M, Levy SE, Mandell DS, Weaver TE, Pinto-Martin J. Sleep behaviors

- and sleep quality in children with autism spectrum disorders. *Sleep*. 2009 Dec;32(12):1566-78.
- Stein MA, Mendelsohn J, Obermeyer WH, Amromin J, Benca R. Sleep and behavior problems in school-aged children. *Pediatrics*. 2001 Apr;107(4):E60.
- Taveras EM, Rifas-Shiman SL, Oken E, Gunderson EP, Gillman MW. Short sleep duration in infancy and risk of childhood overweight. *Arch Pediatr Adolesc Med*. 2008 Apr;162(4):305-11.
- Thomas DA, Poole K, McArdle EK, Goodenough PC, Thompson J, Beardsmore CS, Simpson H. The effect of sleep deprivation on sleep states, breathing events, peripheral chemoresponsiveness and arousal propensity in healthy 3 month old infants. *Eur Respir J*. 1996 May;9(5):932-8.
- Tirosh E, Scher A, Sadeh A, Jaffe M, Lavie P. Sleep characteristics of asthmatics in the first four years of life: a comparative study. *Arch Dis Child*. 1993 Apr;68(4):481-3.
- Touchette E, Petit D, Tremblay RE, Boivin M, Falissard B, Genolini C, Montplaisir JY. Associations between sleep duration patterns and overweight/obesity at age 6. *Sleep*. 2008 Nov;31(11):1507-14.
- Touchette E, Petit D, Tremblay RE, Montplaisir JY. Risk factors and consequences of early childhood dyssomnias: New perspectives. *Sleep Med Rev*. 2009 Oct;13(5):355-61.
- Wang G, Xu G, Liu Z, Lu N, Ma R, Zhang E. Sleep patterns and sleep disturbances among Chinese school-aged children: prevalence and associated factors. *Sleep Med*. 2013 Jan;14(1):45-52.
- Weissbluth M, Brouillette RT, Liu K, Hunt CE. Sleep apnea, sleep duration, and infant temperament. *J Pediatr*. 1982 Aug;101(2):307-10.
- Winsler A, Deutsch A, Vorona RD, Payne PA, Szklo-Coxe M. Sleepless in Fairfax: the difference one more hour of sleep can make for teen hopelessness, suicidal ideation, and substance use. *J Youth Adolesc*. 2015 Feb;44(2):362-78.
- Wong WW, Ortiz CL, Lathan D, Moore LA, Konzelmann KL, Adolph AL, Smith EO, Butte NF. Sleep duration of underserved minority children in a cross-sectional study. *BMC Public Health*. 2013 Jul 12;13:648.
- Yang CK, Kim JK, Patel SR, Lee JH. Age-related changes in sleep/wake patterns among Korean teenagers. *Pediatrics*. 2005 Jan;115(1 Suppl):250-6.
- Yen CF, King BH, Tang TC. The association between short and long nocturnal sleep durations and risky behaviours and the moderating factors in Taiwanese adolescents. *Psychiatry Res*. 2010 Aug 30;179(1):69-74.
- Yen CF, Ko CH, Yen JY, Cheng CP. The multidimensional correlates associated with short nocturnal sleep duration and subjective insomnia among Taiwanese adolescents. *Sleep*. 2008 Nov;31(11):1515-25.
- Alfano CA, Pina AA, Zerr AA, Villalta IK. Pre-sleep arousal and sleep problems of anxiety-disordered youth. *Child Psychiatry Hum Dev*. 2010 Apr;41(2):156-67.
- Arman AR, Ay P, Fis NP, Ersu R, Topuzoglu A, Isik U, Berkem M. Association of sleep duration with socio-economic status and behavioural problems among schoolchildren. *Acta Paediatr*. 2011 Mar;100(3):420-4.
- Aronen ET, Paavonen EJ, Fjällberg M, Soininen M, Törrönen J. Sleep and psychiatric symptoms in school-age children. *J Am Acad Child Adolesc Psychiatry*. 2000 Apr;39(4):502-8.
- Asarnow LD, McGlinchey E, Harvey AG. The effects of bedtime and sleep duration on academic and emotional outcomes in a nationally representative sample of adolescents. *J Adolesc Health*. 2014 Mar;54(3):350-6.
- Astill RG, Van der Heijden KB, Van Ijzendoorn MH, Van Someren EJ. Sleep, cognition, and behavioral problems in school-age children: a century of research meta-analyzed. *Psychol Bull*. 2012 Nov;138(6):1109-38.
- Barnes JC, Meldrum RC. The impact of sleep duration on adolescent development: a genetically informed analysis of identical twin pairs. *J Youth Adolesc*. 2015 Feb;44(2):489-506.
- Baum KT, Desai A, Field J, Miller LE, Rausch J, Beebe DW. Sleep restriction worsens mood and emotion regulation in adolescents. *J Child Psychol Psychiatry*. 2014;55(2):180-90.
- Beebe DW, Ris MD, Kramer ME, Long E, Amin R. The association between sleep disordered breathing, academic grades, and cognitive and behavioral functioning among overweight subjects during middle to late childhood. *Sleep*. 2010 Nov;33(11):1447-56.
- Berger RH, Miller AL, Seifer R, Cares SR, LeBourgeois MK. Acute sleep restriction effects on emotion responses in 30- to 36-month-old children. *J Sleep Res*. 2012 Jun;21(3):235-46.
- Biggs SN, Lushington K, van den Heuvel CJ, Martin AJ, Kennedy JD. Inconsistent sleep schedules and daytime behavioral difficulties in school-aged children. *Sleep Med*. 2011 Sep;12(8):780-6.
- Burt J, Dube L, Thibault L, Gruber R. Sleep and eating in childhood: a potential behavioral mechanism underlying the relationship between poor sleep and obesity. *Sleep Med*. 2014 Jan;15(1):71-5.
- Capaldi li VF, Handwerker K, Richardson E, Stroud LR. Associations between sleep and cortisol responses to stress in children and adolescents: a pilot study. *Behav Sleep Med*. 2005;3(4):177-92.
- Carskadon MA. Sleep in adolescents: the perfect storm. *Pediatr Clin North Am*. 2011 Jun;58(3):637-47.
- Carvalho Bos S, Gomes A, Clemente V, Marques M, Pereira AT, Maia B, Soares MJ, Cabral AS, Macedo A, Gozal D, Azevedo MH. Sleep and behavioral/emotional problems in children: a population-based study. *Sleep Med*. 2009 Jan;10(1):66-74.
- Casoff J, Bhatti JA, Gruber R. The effect of sleep restriction on neurobehavioural functioning in normally developing children and adolescents: insights from the Attention, Behaviour and Sleep Laboratory. *Pathol Biol (Paris)*. 2014 Oct;62(5):319-31.

Mental Health

- Accardo JA, Marcus CL, Leonard MB, Shults J, Meltzer LJ, Elia J. Associations between psychiatric comorbidities and sleep disturbances in children with attention-deficit/hyperactivity disorder. *J Dev Behav Pediatr*. 2012 Feb;33(2):97-105.

- Cortese S, Konofal E, Yateman N, Mouren MC, Lecendreux M. Sleep and alertness in children with attention-deficit/hyperactivity disorder: a systematic review of the literature. *Sleep*. 2006 Apr;29(4):504-11.
- Cortesi F, Giannotti F, Sebastiani T, Vagnoni C, Marioni P. Cosleeping versus solitary sleeping in children with bedtime problems: child emotional problems and parental distress. *Behav Sleep Med*. 2008;6(2):89-105.
- Countryman AJ, Saab PG, Llabre MM, Penedo FJ, McCalla JR, Schneiderman N. Cardiometabolic risk in adolescents: associations with physical activity, fitness, and sleep. *Ann Behav Med*. 2013 Feb;45(1):121-31.
- Dagys N, McGlinchey EL, Talbot LS, Kaplan KA, Dahl RE, Harvey AG. Double trouble? The effects of sleep deprivation and chronotype on adolescent affect. *J Child Psychol Psychiatry*. 2012 Jun;53(6):660-7.
- Dahl RE, Lewin DS. Pathways to adolescent health sleep regulation and behavior. *J Adolesc Health*. 2002 Dec;31(6 Suppl):175-84.
- Dahl RE, Ryan ND, Matty MK, Birmaher B, al-Shabbout M, Williamson DE, Kupfer DJ. Sleep onset abnormalities in depressed adolescents. *Biol Psychiatry*. 1996 Mar 15;39(6):400-10.
- Daniel LC, Boergers J, Kopel SJ, Koinis-Mitchell D. Missed sleep and asthma morbidity in urban children. *Ann Allergy Asthma Immunol*. 2012 Jul;109(1):41-6.
- Do YK, Shin E, Bautista MA, Foo K. The associations between self-reported sleep duration and adolescent health outcomes: what is the role of time spent on Internet use? *Sleep Med*. 2013 Feb;14(2):195-200.
- Doane LD, Thurston EC. Associations among sleep, daily experiences, and loneliness in adolescence: evidence of moderating and bidirectional pathways. *J Adolesc*. 2014 Feb;37(2):145-54.
- El-Sheikh M, Buckhalt JA, Mark Cummings E, Keller P. Sleep disruptions and emotional insecurity are pathways of risk for children. *J Child Psychol Psychiatry*. 2007 Jan;48(1):88-96.
- Ertan P, Yilmaz O, Caglayan M, Sogut A, Aslan S, Yuksel H. Relationship of sleep quality and quality of life in children with monosymptomatic enuresis. *Child Care Health Dev*. 2009 Jul;35(4):469-74.
- Fallone G, Acebo C, Seifer R, Carskadon MA. Experimental restriction of sleep opportunity in children: effects on teacher ratings. *Sleep*. 2005 Dec;28(12):1561-7.
- Fallone G, Acebo C, Arnedt JT, Seifer R, Carskadon MA. Effects of acute sleep restriction on behavior, sustained attention, and response inhibition in children. *Percept Mot Skills*. 2001 Aug;93(1):213-29.
- Fredriksen K, Rhodes J, Reddy R, Way N. Sleepless in Chicago: tracking the effects of adolescent sleep loss during the middle school years. *Child Dev*. 2004 Jan-Feb;75(1):84-95.
- Gau SS, Shang CY, Merikangas KR, Chiu YN, Soong WT, Cheng AT. Association between morningness-eveningness and behavioral/emotional problems among adolescents. *J Biol Rhythms*. 2007 Jun;22(3):268-74.
- Giannotti F, Cortesi F, Cerquiglini A, Miraglia D, Vagnoni C, Sebastiani T, Bernabei P. An investigation of sleep characteristics, EEG abnormalities and epilepsy in developmentally regressed and non-regressed children with autism. *J Autism Dev Disord*. 2008 Nov;38(10):1888-97.
- Gruber R, Cassoff J, Frenette S, Wiebe S, Carrier J. Impact of sleep extension and restriction on children's emotional lability and impulsivity. *Pediatrics*. 2012 Nov;130(5):e1155-61.
- Gruber R, Wiebe S, Montecalvo L, Brunetti B, Amsel R, Carrier J. Impact of sleep restriction on neurobehavioral functioning of children with attention deficit hyperactivity disorder. *Sleep*. 2011 Mar 1;34(3):315-23.
- Gruber R, Xi T, Frenette S, Robert M, Vannasinh P, Carrier J. Sleep disturbances in prepubertal children with attention deficit hyperactivity disorder: a home polysomnography study. *Sleep*. 2009 Mar;32(3):343-50.
- Gruber R, Sadeh A, Raviv A. Instability of sleep patterns in children with attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry*. 2000 Apr;39(4):495-501.
- Hansen M, Janssen I, Schiff A, Zee PC, Dubocovich ML. The impact of school daily schedule on adolescent sleep. *Pediatrics*. 2005 Jun;115(6):1555-61.
- Hiscock H, Sciberras E, Mensah F, Gerner B, Efron D, Khano S, Oberklaid F. Impact of a behavioural sleep intervention on symptoms and sleep in children with attention deficit hyperactivity disorder, and parental mental health: randomised controlled trial. *BMJ*. 2015 Jan 20;350:h68.
- Hodge D, Carollo TM, Lewin M, Hoffman CD, Sweeney DP. Sleep patterns in children with and without autism spectrum disorders: developmental comparisons. *Res Dev Disabil*. 2014 Jul;35(7):1631-8.
- Holley S, Hill CM, Stevenson J. An hour less sleep is a risk factor for childhood conduct problems. *Child Care Health Dev*. 2011 Jul;37(4):563-70.
- Honomichl RD, Goodlin-Jones BL, Burnham M, Gaylor E, Anders TF. Sleep patterns of children with pervasive developmental disorders. *J Autism Dev Disord*. 2002 Dec;32(6):553-61.
- Humphreys JS, Gringras P, Blair PS, Scott N, Henderson J, Fleming PJ, Emond AM. Sleep patterns in children with autistic spectrum disorders: a prospective cohort study. *Arch Dis Child*. 2014 Feb;99(2):114-8.
- Kang SG, Lee YJ, Kim SJ, Lim W, Lee HJ, Park YM, Cho IH, Cho SJ, Hong JP. Weekend catch-up sleep is independently associated with suicide attempts and self-injury in Korean adolescents. *Compr Psychiatry*. 2014 Feb;55(2):319-25.
- Keller PS, El-Sheikh M, Buckhalt JA. Children's attachment to parents and their academic functioning: sleep disruptions as moderators of effects. *J Dev Behav Pediatr*. 2008 Dec;29(6):441-9.
- Kelly RJ, El-Sheikh M. Reciprocal relations between children's sleep and their adjustment over time. *Dev Psychol*. 2014 Apr;50(4):1137-47.
- Kim HS, Ham OK, Kim JW, Park JY. Association between sleep duration and psychological health in overweight and obese children in Korea. *Nurs Health Sci*. 2012 Jun;14(2):238-43.
- Komada Y, Abe T, Okajima I, Asaoka S, Matsuura N, Usui A, Shirakawa S, Inoue Y. Short sleep duration and

- irregular bedtime are associated with increased behavioral problems among Japanese preschool-age children. *Tohoku J Exp Med.* 2011;224(2):127-36.
- Kouros CD, El-Sheikh M. Daily mood and sleep: reciprocal relations and links with adjustment problems. *J Sleep Res.* 2015 Feb;24(1):24-31.
- Kwon JA, Lee M, Yoo KB, Park EC. Does the duration and time of sleep increase the risk of allergic rhinitis? Results of the 6-year nationwide Korea youth risk behavior web-based survey. *PLoS One.* 2013;8(8):e72507.
- Lee YJ, Cho SJ, Cho IH, Kim SJ. Insufficient sleep and suicidality in adolescents. *Sleep.* 2012 Apr 1;35(4):455-60.
- Lehto JE, Uusitalo-Malmivaara L. Sleep-related factors: associations with poor attention and depressive symptoms. *Child Care Health Dev.* 2014 May;40(3):419-25.
- Lewandowski AS, Palermo TM, De la Motte S, Fu R. Temporal daily associations between pain and sleep in adolescents with chronic pain versus healthy adolescents. *Pain.* 2010 Oct;151(1):220-5.
- Lin WH, Yi CC. Unhealthy sleep practices, conduct problems, and daytime functioning during adolescence. *J Youth Adolesc.* 2015 Feb;44(2):431-46.
- Liu X, Forbes EE, Ryan ND, Rofey D, Hannon TS, Dahl RE. Rapid eye movement sleep in relation to overweight in children and adolescents. *Arch Gen Psychiatry.* 2008 Aug;65(8):924-32.
- Liu X, Hubbard JA, Fabes RA, Adam JB. Sleep disturbances and correlates of children with autism spectrum disorders. *Child Psychiatry Hum Dev.* 2006 Winter;37(2):179-91.
- Liu X. Sleep and adolescent suicidal behavior. *Sleep.* 2004 Nov 1;27(7):1351-8.
- Liu X, Zhou H. Sleep duration, insomnia and behavioral problems among Chinese adolescents. *Psychiatry Res.* 2002 Aug 5;111(1):75-85.
- Lytle LA, Pasch KE, Farbakhsh K. The relationship between sleep and weight in a sample of adolescents. *Obesity (Silver Spring).* 2011 Feb;19(2):324-31.
- Magee CA, Gordon R, Caputi P. Distinct developmental trends in sleep duration during early childhood. *Pediatrics.* 2014 Jun;133(6):e1561-7.
- Malow BA, Marzec ML, McGrew SG, Wang L, Henderson LM, Stone WL. Characterizing sleep in children with autism spectrum disorders: a multidimensional approach. *Sleep.* 2006 Dec;29(12):1563-71.
- Maski KP, Kothare SV. Sleep deprivation and neurobehavioral functioning in children. *Int J Psychophysiol.* 2013 Aug;89(2):259-64.
- Matamura M, Tochigi M, Usami S, Yonehara H, Fukushima M, Nishida A, Togo F, Sasaki T. Associations between sleep habits and mental health status and suicidality in a longitudinal survey of monozygotic twin adolescents. *J Sleep Res.* 2014 Jun;23(3):290-4.
- Matsushita M, Koyama A, Ushijima H, Mikami A, Katsumata Y, Kikuchi Y, Ichimi N, Jono T, Fujise N, Ikeda M. Sleep duration and its association with sleepiness and depression in ronin-sei" preparatory school students." *Asian J Psychiatr.* 2014 Jun;9:61-6.
- McDermott BM, Mamun AA, Najman JM, Williams GM, O'callaghan MJ, Bor W. Preschool children perceived by mothers as irregular eaters: physical and psychosocial predictors from a birth cohort study. *J Dev Behav Pediatr.* 2008 Jun;29(3):197-205.
- Meltzer LJ, Mindell JA. Sleep and sleep disorders in children and adolescents. *Psychiatr Clin North Am.* 2006 Dec;29(4):1059-76
- Mercier L, Pivik RT, Busby K. Sleep patterns in reading disabled children. *Sleep.* 1993 Apr;16(3):207-15.
- Moore M, Kirchner HL, Drotar D, Johnson N, Rosen C, Redline S. Correlates of adolescent sleep time and variability in sleep time: the role of individual and health related characteristics. *Sleep Med.* 2011 Mar;12(3):239-45.
- Moore M, Kirchner HL, Drotar D, Johnson N, Rosen C, Ancoli-Israel S, Redline S. Relationships among sleepiness, sleep time, and psychological functioning in adolescents. *J Pediatr Psychol.* 2009 Nov-Dec;34(10):1175-83.
- Moseley L, Gradisar M. Evaluation of a school-based intervention for adolescent sleep problems. *Sleep.* 2009 Mar;32(3):334-41.
- Owens J. Classification and epidemiology of childhood sleep disorders. *Prim Care.* 2008 Sep;35(3):533-46, vii.
- Owens JA, Belon K, Moss P. Impact of delaying school start time on adolescent sleep, mood, and behavior. *Arch Pediatr Adolesc Med.* 2010 Jul;164(7):608-14.
- Owens JA, Fernando S, Mc Guinn M. Sleep disturbance and injury risk in young children. *Behav Sleep Med.* 2005;3(1):18-31.
- Owens JA, Maxim R, Nobile C, McGuinn M, Msall M. Parental and self-report of sleep in children with attention-deficit/hyperactivity disorder. *Arch Pediatr Adolesc Med.* 2000 Jun;154(6):549-55.
- Paavonen EJ, Räikkönen K, Lahti J, Komsu N, Heinonen K, Pesonen AK, Järvenpää AL, Strandberg T, Kajantie E, Porkka-Heiskanen T. Short sleep duration and behavioral symptoms of attention-deficit/hyperactivity disorder in healthy 7- to 8-year-old children. *Pediatrics.* 2009 May;123(5):e857-64.
- Paavonen EJ, Porkka-Heiskanen T, Lahikainen AR. Sleep quality, duration and behavioral symptoms among 5-6-year-old children. *Eur Child Adolesc Psychiatry.* 2009 Dec;18(12):747-54.
- Paavonen EJ, Vehkalahti K, Vanhala R, von Wendt L, Nieminen-von Wendt T, Aronen ET. Sleep in children with Asperger syndrome. *J Autism Dev Disord.* 2008 Jan;38(1):41-51
- Paavonen EJ, Nieminen-von Wendt T, Vanhala R, Aronen ET, von Wendt L. Effectiveness of melatonin in the treatment of sleep disturbances in children with Asperger disorder. *J Child Adolesc Psychopharmacol.* 2003 Spring;13(1):83-95.
- Paciência I, Barros H, Araújo J, Ramos E. Association between sleep duration and blood pressure in adolescents. *Hypertens Res.* 2013 Aug;36(8):747-52.
- Pantic I, Damjanovic A, Todorovic J, Topalovic D, Bojovic-Jovic D, Ristic S, Pantic S. Association between online social networking and depression in high school students: behavioral physiology viewpoint. *Psychiatr Danub.* 2012 Mar;24(1):90-3.

- Pasch KE, Laska MN, Lytle LA, Moe SG. Adolescent sleep, risk behaviors, and depressive symptoms: are they linked? *Am J Health Behav.* 2010 Mar-Apr;34(2):237-48.
- Perez-Chada D, Perez-Lloret S, Videla AJ, Cardinali D, Bergna MA, Fernández-Acquier M, Larrateguy L, Zabert GE, Drake C. Sleep disordered breathing and daytime sleepiness are associated with poor academic performance in teenagers. A study using the Pediatric Daytime Sleepiness Scale (PDSS). *Sleep.* 2007 Dec;30(12):1698-703.
- Perkinson-Gloor N, Lemola S, Grob A. Sleep duration, positive attitude toward life, and academic achievement: the role of daytime tiredness, behavioral persistence, and school start times. *J Adolesc.* 2013 Apr;36(2):311-8.
- Roberts RE, Duong HT. The prospective association between sleep deprivation and depression among adolescents. *Sleep.* 2014 Feb 1;37(2):239-44.
- Sadeh A, Gruber R, Raviv A. Sleep, neurobehavioral functioning, and behavior problems in school-age children. *Child Dev.* 2002 Mar-Apr;73(2):405-17.
- Scharf RJ, Demmer RT, Silver EJ, Stein RE. Nighttime sleep duration and externalizing behaviors of preschool children. *J Dev Behav Pediatr.* 2013 Jul-Aug;34(6):384-91.
- Schreck KA, Mulick JA, Smith AF. Sleep problems as possible predictors of intensified symptoms of autism. *Res Dev Disabil.* 2004 Jan-Feb;25(1):57-66.
- Scott N, Blair PS, Emond AM, Fleming PJ, Humphreys JS, Henderson J, Gringras P. Sleep patterns in children with ADHD: a population-based cohort study from birth to 11 years. *J Sleep Res.* 2013 Apr;22(2):121-8.
- Segura-Jiménez V, Carbonell-Baeza A, Keating XD, Ruiz JR, Castro-Piñero J. Association of sleep patterns with psychological positive health and health complaints in children and adolescents. *Qual Life Res.* 2015 Apr;24(4):885-95.
- Short MA, Gradisar M, Lack LC, Wright HR, Dohnt H. The sleep patterns and well-being of Australian adolescents. *J Adolesc.* 2013 Feb;36(1):103-10.
- Short MA, Gradisar M, Lack LC, Wright HR. The impact of sleep on adolescent depressed mood, alertness and academic performance. *J Adolesc.* 2013 Dec;36(6):1025-33.
- Shur-Fen Gau S. Prevalence of sleep problems and their association with inattention/hyperactivity among children aged 6-15 in Taiwan. *J Sleep Res.* 2006 Dec;15(4):403-14. Erratum in: *J Sleep Res.* 2010 Jun;19(2):379. Shur-Fen, Gau Susan [corrected to Shur-Fen Gau, Susan].
- Silva GE, Goodwin JL, Parthasarathy S, Sherrill DL, Vana KD, Drescher AA, Quan SF. Longitudinal association between short sleep, body weight, and emotional and learning problems in Hispanic and Caucasian children. *Sleep.* 2011 Sep 1;34(9):1197-205.
- Sivertsen B, Harvey AG, Reichborn-Kjennerud T, Torgersen L, Ystrom E, Hysing M. Later emotional and behavioral problems associated with sleep problems in toddlers: a longitudinal study. *JAMA Pediatr.* 2015 Jun;169(6):575-82.
- Sivertsen B, Harvey AG, Lundervold AJ, Hysing M. Sleep problems and depression in adolescence: results from a large population-based study of Norwegian adolescents aged 16-18 years. *Eur Child Adolesc Psychiatry.* 2014 Aug;23(8):681-9.
- Smaldone A, Honig JC, Byrne MW. Sleepless in America: inadequate sleep and relationships to health and well-being of our nation's children. *Pediatrics.* 2007 Feb;119 Suppl 1:S29-37.
- Souders MC, Mason TB, Valladares O, Bucan M, Levy SE, Mandell DS, Weaver TE, Pinto-Martin J. Sleep behaviors and sleep quality in children with autism spectrum disorders. *Sleep.* 2009 Dec;32(12):1566-78.
- Spruyt K, Gozal D. Sleep disturbances in children with attention-deficit/hyperactivity disorder. *Expert Rev Neurother.* 2011 Apr;11(4):565-77.
- Stein MA, Mendelsohn J, Obermeyer WH, Amromin J, Benca R. Sleep and behavior problems in school-aged children. *Pediatrics.* 2001 Apr;107(4):E60.
- Suzuki H, Kaneita Y, Osaki Y, Minowa M, Kanda H, Suzuki K, Wada K, Hayashi K, Tanihata T, Ohida T. Clarification of the factor structure of the 12-item General Health Questionnaire among Japanese adolescents and associated sleep status. *Psychiatry Res.* 2011 Jun 30;188(1):138-46.
- Tininenko JR, Fisher PA, Bruce J, Pears KC. Associations between sleep and inattentive/hyperactive problem behavior among foster and community children. *J Dev Behav Pediatr.* 2010 Oct;31(8):668-74.
- Touchette E, Petit D, Tremblay RE, Montplaisir JY. Risk factors and consequences of early childhood dyssomnias: New perspectives. *Sleep Med Rev.* 2009 Oct;13(5):355-61.
- Touchette E, Côté SM, Petit D, Liu X, Boivin M, Falissard B, Tremblay RE, Montplaisir JY. Short nighttime sleep-duration and hyperactivity trajectories in early childhood. *Pediatrics.* 2009 Nov;124(5):e985-93.
- Vazsonyi AT, Harris C, Terveer AM, Pagava K, Phagava H, Michaud PA. Parallel mediation effects by sleep on the parental warmth-problem behavior links: evidence from national probability samples of Georgian and Swiss adolescents. *J Youth Adolesc.* 2015 Feb;44(2):331-45.
- Vriend JL, Davidson FD, Corkum PV, Rusak B, Chambers CT, McLaughlin EN. Manipulating sleep duration alters emotional functioning and cognitive performance in children. *J Pediatr Psychol.* 2013 Nov;38(10):1058-69.
- Ward TM, Gay C, Alkon A, Anders TF, Lee KA. Nocturnal sleep and daytime nap behaviors in relation to salivary cortisol levels and temperament in preschool-age children attending child care. *Biol Res Nurs.* 2008 Jan;9(3):244-53.
- Wiggs L, Stores G. Behavioural treatment for sleep problems in children with severe learning disabilities and challenging daytime behaviour: effect on daytime behaviour. *J Child Psychol Psychiatry.* 1999 May;40(4):627-35.
- Willoughby MT, Angold A, Egger HL. Parent-reported attention-deficit/hyperactivity disorder symptomatology and sleep problems in a preschool-age pediatric clinic sample. *J Am Acad Child Adolesc Psychiatry.* 2008 Sep;47(9):1086-94.
- Winsler A, Deutsch A, Vorona RD, Payne PA, Szklo-Coxe M. Sleepless in Fairfax: the difference one more hour of sleep can make for teen hopelessness, suicidal ideation,

and substance use. *J Youth Adolesc.* 2015 Feb;44(2):362-78.

Yen CF, King BH, Tang TC. The association between short and long nocturnal sleep durations and risky behaviours and the moderating factors in Taiwanese adolescents. *Psychiatry Res.* 2010 Aug 30;179(1):69-74.

Yen CF, Ko CH, Yen JY, Cheng CP. The multidimensional correlates associated with short nocturnal sleep duration and subjective insomnia among Taiwanese adolescents. *Sleep.* 2008 Nov;31(11):1515-25.

Yip T. The effects of ethnic/racial discrimination and sleep quality on depressive symptoms and self-esteem trajectories among diverse adolescents. *J Youth Adolesc.* 2015 Feb;44(2):419-30.

Yokomaku A, Misao K, Omoto F, Yamagishi R, Tanaka K, Takada K, Kohyama J. A study of the association between sleep habits and problematic behaviors in preschool children. *Chronobiol Int.* 2008 Jul;25(4):549-64.

Metabolic Health

Agras WS, Hammer LD, McNicholas F, Kraemer HC. Risk factors for childhood overweight: a prospective study from birth to 9.5 years. *J Pediatr.* 2004 Jul;145(1):20-5. Erratum in: *J Pediatr.* 2004 Sep;145(3):424.

Al-Disi D, Al-Daghri N, Khanam L, Al-Othman A, Al-Saif M, Sabico S, Chrousos G. Subjective sleep duration and quality influence diet composition and circulating adipocytokines and ghrelin levels in teen-age girls. *Endocr J.* 2010;57(10):915-23.

Al-Hazzaa HM, Musaiger AO, Abahussain NA, Al-Sobayel HI, Qahwaji DM. Lifestyle correlates of self-reported sleep duration among Saudi adolescents: a multicentre school-based cross-sectional study. *Child Care Health Dev.* 2014 Jul;40(4):533-42.

Altenburg TM, Chinapaw MJ, van der Knaap ET, Brug J, Manios Y, Singh AS. Longer sleep--slimmer kids: the ENERGY-project. *PLoS One.* 2013;8(3):e59522.

Androutsos O, Moschonis G, Mavrogianni C, Roma-Giannikou E, Chrousos GP, Kanaka-Gantenbein C, Manios Y. Identification of lifestyle patterns, including sleep deprivation, associated with insulin resistance in children: the Healthy Growth Study. *Eur J Clin Nutr.* 2014 Mar;68(3):344-9.

Appelhans BM, Fitzpatrick SL, Li H, Cail V, Waring ME, Schneider KL, Whited MC, Busch AM, Pagoto SL. The home environment and childhood obesity in low-income households: indirect effects via sleep duration and screen time. *BMC Public Health.* 2014 Nov 9;14:1160.

Arora T, Taheri S. Associations among late chronotype, body mass index and dietary behaviors in young adolescents. *Int J Obes (Lond).* 2015 Jan;39(1):39-44.

Arora T, Hussain S, Hubert Lam KB, Lily Yao G, Neil Thomas G, Taheri S. Exploring the complex pathways among specific types of technology, self-reported sleep duration and body mass index in UK adolescents. *Int J Obes (Lond).* 2013 Sep;37(9):1254-60.

Arora T, Hosseini-Araghi M, Bishop J, Yao GL, Thomas GN, Taheri S. The complexity of obesity in U.K. adolescents: relationships with quantity and type of technology, sleep

duration and quality, academic performance and aspiration. *Pediatr Obes.* 2013 Oct;8(5):358-66.

Azadbakht L, Kelishadi R, Khodarahmi M, Qorbani M, Heshmat R, Motlagh ME, Taslimi M, Ardalan G. The association of sleep duration and cardiometabolic risk factors in a national sample of children and adolescents: the CASPIAN III study. *Nutrition.* 2013 Sep;29(9):1133-41.

Ball GD, Lenk JM, Barbarich BN, Plotnikoff RC, Fishburne GJ, Mackenzie KA, Willows ND. Overweight children and adolescents referred for weight management: are they meeting lifestyle behaviour recommendations? *Appl Physiol Nutr Metab.* 2008 Oct;33(5):936-45.

Beebe DW, Ris MD, Kramer ME, Long E, Amin R. The association between sleep disordered breathing, academic grades, and cognitive and behavioral functioning among overweight subjects during middle to late childhood. *Sleep.* 2010 Nov;33(11):1447-56.

Beebe DW, Lewin D, Zeller M, McCabe M, MacLeod K, Daniels SR, Amin R. Sleep in overweight adolescents: shorter sleep, poorer sleep quality, sleepiness, and sleep-disordered breathing. *J Pediatr Psychol.* 2007 Jan-Feb;32(1):69-79.

Bell JF, Zimmerman FJ. Shortened nighttime sleep duration in early life and subsequent childhood obesity. *Arch Pediatr Adolesc Med.* 2010 Sep;164(9):840-5. Erratum in: *Arch Pediatr Adolesc Med.* 2010 Nov;164(11):1070.

Bibiloni Mdel M, Martinez E, Llull R, Juarez MD, Pons A, Tur JA. Prevalence and risk factors for obesity in Balearic Islands adolescents. *Br J Nutr.* 2010 Jan;103(1):99-106.

Bonuck K, Chervin RD, Howe LD. Sleep-disordered breathing, sleep duration, and childhood overweight: a longitudinal cohort study. *J Pediatr.* 2015 Mar;166(3):632-9.

Brandalize M, Pereira RF, Leite N, Filho GL, Louzada FM. Effect of morning school schedule on sleep and anthropometric variables in adolescents: a follow-up study. *Chronobiol Int.* 2011 Nov;28(9):779-85

Burt J, Dube L, Thibault L, Gruber R. Sleep and eating in childhood: a potential behavioral mechanism underlying the relationship between poor sleep and obesity. *Sleep Med.* 2014 Jan;15(1):71-5.

Börnhorst C, Hense S, Ahrens W, Hebestreit A, Reisch L, Barba G, von Kries R, Bayer O; IDEFICS Consortium. From sleep duration to childhood obesity--what are the pathways? *Eur J Pediatr.* 2012 Jul;171(7):1029-38.

Calamaro CJ, Park S, Mason TB, Marcus CL, Weaver TE, Pack A, Ratcliffe SJ. Shortened sleep duration does not predict obesity in adolescents. *J Sleep Res.* 2010 Dec;19(4):559-66.

Carrillo-Larco RM, Bernabé-Ortiz A, Miranda JJ. Short sleep duration and childhood obesity: cross-sectional analysis in Peru and patterns in four developing countries. *PLoS One.* 2014;9(11):e112433.

Casazza K, Hanks LJ, Fernandez JR. Shorter sleep may be a risk factor for impaired bone mass accrual in childhood. *J Clin Densitom.* 2011 Oct-Dec;14(4):453-7.

Casoff J, Bhatti JA, Gruber R. The effect of sleep restriction on neurobehavioural functioning in normally developing children and adolescents: insights from the Attention,

- Behaviour and Sleep Laboratory. *Pathol Biol (Paris)*. 2014 Oct;62(5):319-31.
- Cespedes EM, Rifas-Shiman SL, Redline S, Gillman MW, Peña MM, Taveras EM. Longitudinal associations of sleep curtailment with metabolic risk in mid-childhood. *Obesity (Silver Spring)*. 2014 Dec;22(12):2586-92.
- Chamorro R, Algarín C, Garrido M, Causa L, Held C, Lozoff B, Peirano P. Night time sleep macrostructure is altered in otherwise healthy 10-year-old overweight children. *Int J Obes (Lond)*. 2014 Aug;38(8):1120-5.
- Chaput JP, Lambert M, Gray-Donald K, McGrath JJ, Tremblay MS, O'Loughlin J, Tremblay A. Short sleep duration is independently associated with overweight and obesity in Quebec children. *Can J Public Health*. 2011 Sep-Oct;102(5):369-74.
- Chaput JP, Brunet M, Tremblay A. Relationship between short sleeping hours and childhood overweight/obesity: results from the 'Québec en Forme' Project. *Int J Obes (Lond)*. 2006 Jul;30(7):1080-5.
- Chen X, Beydoun MA, Wang Y. Is sleep duration associated with childhood obesity? A systematic review and meta-analysis. *Obesity (Silver Spring)*. 2008 Feb;16(2):265-74.
- Cicek B, Ozturk A, Mazicioglu MM, Elmali F, Turp N, Kurtoglu S. The risk analysis of arm fat area in Turkish children and adolescents. *Ann Hum Biol*. 2009 Jan-Feb;36(1):28-37.
- Countryman AJ, Saab PG, Llabre MM, Penedo FJ, McCalla JR, Schneiderman N. Cardiometabolic risk in adolescents: associations with physical activity, fitness, and sleep. *Ann Behav Med*. 2013 Feb;45(1):121-31.
- D'Aniello R, Troisi J, D'Amico O, Sangermano M, Massa G, Moccaldò A, Pierri L, Poeta M, Vajro P. Emerging pathomechanisms involved in obesity. *J Pediatr Gastroenterol Nutr*. 2015 Jan;60(1):113-9.
- Danielsen YS, Pallesen S, Stormark KM, Nordhus IH, Bjorvatn B. The relationship between school day sleep duration and body mass index in Norwegian children (aged 10-12). *Int J Pediatr Obes*. 2010 May 3;5(3):214-20.
- de Jong E, Stocks T, Visscher TL, HiraSing RA, Seidell JC, Renders CM. Association between sleep duration and overweight: the importance of parenting. *Int J Obes (Lond)*. 2012 Oct;36(10):1278-84.
- Dev DA, McBride BA, Fiese BH, Jones BL, Cho H; Behalf Of The Strong Kids Research Team. Risk factors for overweight/obesity in preschool children: an ecological approach. *Child Obes*. 2013 Oct;9(5):399-408.
- Diethelm K, Bolzenius K, Cheng G, Remer T, Buyken AE. Longitudinal associations between reported sleep duration in early childhood and the development of body mass index, fat mass index and fat free mass index until age 7. *Int J Pediatr Obes*. 2011 Jun;6(2-2):e114-23.
- Do YK, Shin E, Bautista MA, Foo K. The associations between self-reported sleep duration and adolescent health outcomes: what is the role of time spent on Internet use? *Sleep Med*. 2013 Feb;14(2):195-200.
- Drescher AA, Goodwin JL, Silva GE, Quan SF. Caffeine and screen time in adolescence: associations with short sleep and obesity. *J Clin Sleep Med*. 2011 Aug 15;7(4):337-42.
- Duncan JS, Schofield G, Duncan EK, Rush EC. Risk factors for excess body fatness in New Zealand children. *Asia Pac J Clin Nutr*. 2008;17(1):138-47.
- Eisenmann JC, Ekkekakis P, Holmes M. Sleep duration and overweight among Australian children and adolescents. *Acta Paediatr*. 2006 Aug;95(8):956-63.
- Ekstedt M, Nyberg G, Ingre M, Ekblom ö, Marcus C. Sleep, physical activity and BMI in six to ten-year-old children measured by accelerometry: a cross-sectional study. *Int J Behav Nutr Phys Act*. 2013 Jun 22;10:82.
- Fatima Y, Doi SA, Mamun AA. Longitudinal impact of sleep on overweight and obesity in children and adolescents: a systematic review and bias-adjusted meta-analysis. *Obes Rev*. 2015 Feb;16(2):137-49.
- Firouzi S, Poh BK, Ismail MN, Sadeghilar A. Sleep habits, food intake, and physical activity levels in normal and overweight and obese Malaysian children. *Obes Res Clin Pract*. 2014 Jan-Feb;8(1):e70-8.
- Fisher A, McDonald L, van Jaarsveld CH, Llewellyn C, Fildes A, Schrempft S, Wardle J. Sleep and energy intake in early childhood. *Int J Obes (Lond)*. 2014 Jul;38(7):926-9.
- Flint J, Kothare SV, Zihlif M, Suarez E, Adams R, Legido A, De Luca F. Association between inadequate sleep and insulin resistance in obese children. *J Pediatr*. 2007 Apr;150(4):364-9.
- Garaulet M, Ortega FB, Ruiz JR, Rey-López JP, Béghin L, Manios Y, Cuenca-García M, Plada M, Diethelm K, Kafatos A, Molnár D, Al-Tahan J, Moreno LA. Short sleep duration is associated with increased obesity markers in European adolescents: effect of physical activity and dietary habits. The HELENA study. *Int J Obes (Lond)*. 2011 Oct;35(10):1308-17.
- Golley RK, Maher CA, Matricciani L, Olds TS. Sleep duration or bedtime? Exploring the association between sleep timing behaviour, diet and BMI in children and adolescents. *Int J Obes (Lond)*. 2013 Apr;37(4):546-51.
- Gozal D, Kheirandish-Gozal L. Childhood obesity and sleep: relatives, partners, or both?--a critical perspective on the evidence. *Ann N Y Acad Sci*. 2012 Aug;1264:135-41.
- Graef DM, Janicke DM, McCrae CS. Sleep patterns of a primarily obese sample of treatment-seeking children. *J Clin Sleep Med*. 2014 Oct 15;10(10):1111-7.
- Guo X, Zheng L, Li Y, Yu S, Sun G, Yang H, Zhou X, Zhang X, Sun Z, Sun Y. Differences in lifestyle behaviors, dietary habits, and familial factors among normal-weight, overweight, and obese Chinese children and adolescents. *Int J Behav Nutr Phys Act*. 2012 Oct 2;9:120.
- Haines J, McDonald J, O'Brien A, Sherry B, Bottino CJ, Schmidt ME, Taveras EM. Healthy Habits, Happy Homes: randomized trial to improve household routines for obesity prevention among preschool-aged children. *JAMA Pediatr*. 2013 Nov;167(11):1072-9.
- Hart CN, Carskadon MA, Considine RV, Fava JL, Lawton J, Raynor HA, Jelalian E, Owens J, Wing R. Changes in children's sleep duration on food intake, weight, and leptin. *Pediatrics*. 2013 Dec;132(6):e1473-80.
- Hart CN, Cairns A, Jelalian E. Sleep and obesity in children and adolescents. *Pediatr Clin North Am*. 2011 Jun;58(3):715-33.

- Hart CN, Jelalian E. Shortened sleep duration is associated with pediatric overweight. *Behav Sleep Med.* 2008;6(4):251-67.
- Hassan F, Davis MM, Chervin RD. No independent association between insufficient sleep and childhood obesity in the National Survey of Children's Health. *J Clin Sleep Med.* 2011 Apr 15;7(2):153-7.
- Hense S, Pohlabein H, De Henauw S, Eiben G, Molnar D, Moreno LA, Barba G, Hadjigeorgiou C, Veidebaum T, Ahrens W. Sleep duration and overweight in European children: is the association modified by geographic region? *Sleep.* 2011 Jul 1;34(7):885-90.
- Hiscock H, Scalzo K, Canterford L, Wake M. Sleep duration and body mass index in 0-7-year olds. *Arch Dis Child.* 2011 Aug;96(8):735-9.
- Hjorth MF, Chaput JP, Ritz C, Dalskov SM, Andersen R, Astrup A, Tetens I, Michaelsen KF, Sjödin A. Fatness predicts decreased physical activity and increased sedentary time, but not vice versa: support from a longitudinal study in 8- to 11-year-old children. *Int J Obes (Lond).* 2014 Jul;38(7):959-65.
- Hjorth MF, Quist JS, Andersen R, Michaelsen KF, Tetens I, Astrup A, Chaput JP, Sjödin A. Change in sleep duration and proposed dietary risk factors for obesity in Danish school children. *Pediatr Obes.* 2014 Dec;9(6):e156-9.
- Iglayreger HB, Peterson MD, Liu D, Parker CA, Woolford SJ, Sallinen Gafka BJ, Hassan F, Gordon PM. Sleep duration predicts cardiometabolic risk in obese adolescents. *J Pediatr.* 2014 May;164(5):1085-1090.e1.
- Jarrin DC, McGrath JJ, Poirier P, Quality Cohort Collaborative Group. Autonomic dysfunction: a possible pathophysiological pathway underlying the association between sleep and obesity in children at-risk for obesity. *J Youth Adolesc.* 2015 Feb;44(2):285-97.
- Jarrin DC, McGrath JJ, Drake CL. Beyond sleep duration: distinct sleep dimensions are associated with obesity in children and adolescents. *Int J Obes (Lond).* 2013 Apr;37(4):552-8.
- Javaheri S, Storfer-Isser A, Rosen CL, Redline S. Association of short and long sleep durations with insulin sensitivity in adolescents. *J Pediatr.* 2011 Apr;158(4):617-23.
- Jenni OG, Molinari L, Caffisch JA, Largo RH. Sleep duration from ages 1 to 10 years: variability and stability in comparison with growth. *Pediatrics.* 2007 Oct;120(4):e769-76.
- Jiang F, Zhu S, Yan C, Jin X, Bandla H, Shen X. Sleep and obesity in preschool children. *J Pediatr.* 2009 Jun;154(6):814-8.
- Jiang YR, Spruyt K, Chen WJ, Shen XM, Jiang F. Somatic growth of lean children: the potential role of sleep. *World J Pediatr.* 2014 Aug;10(3):245-50.
- Jones CH, Pollard TM, Summerbell CD, Ball H. Could parental rules play a role in the association between short sleep and obesity in young children? *J Biosoc Sci.* 2014 May;46(3):405-18.
- Kim HS, Ham OK, Kim JW, Park JY. Association between sleep duration and psychological health in overweight and obese children in Korea. *Nurs Health Sci.* 2012 Jun;14(2):238-43.
- Kjeldsen JS, Hjorth MF, Andersen R, Michaelsen KF, Tetens I, Astrup A, Chaput JP, Sjödin A. Short sleep duration and large variability in sleep duration are independently associated with dietary risk factors for obesity in Danish school children. *Int J Obes (Lond).* 2014 Jan;38(1):32-9.
- Kleiser C, Schaffrath Rosario A, Mensink GB, Prinz-Langenohl R, Kurth BM. Potential determinants of obesity among children and adolescents in Germany: results from the cross-sectional KiGGS Study. *BMC Public Health.* 2009 Feb 2;9:46.
- Klingenberg L, Christensen LB, Hjorth MF, Zangenberg S, Chaput JP, Sjödin A, Mølgaard C, Michaelsen KF. No relation between sleep duration and adiposity indicators in 9-36 months old children: the SKOT cohort. *Pediatr Obes.* 2013 Feb;8(1):e14-8.
- Klingenberg L, Chaput JP, Holmbäck U, Jennum P, Astrup A, Sjödin A. Sleep restriction is not associated with a positive energy balance in adolescent boys. *Am J Clin Nutr.* 2012 Aug;96(2):240-8.
- Knutson KL. Sex differences in the association between sleep and body mass index in adolescents. *J Pediatr.* 2005 Dec;147(6):830-4.
- Koren D, O'Sullivan KL, Mokhlesi B. Metabolic and glycemic sequelae of sleep disturbances in children and adults. *Curr Diab Rep.* 2015 Jan;15(1):562.
- Koren D, Levitt Katz LE, Brar PC, Gallagher PR, Berkowitz RI, Brooks LJ. Sleep architecture and glucose and insulin homeostasis in obese adolescents. *Diabetes Care.* 2011 Nov;34(11):2442-7.
- Kuriyan R, Bhat S, Thomas T, Vaz M, Kurpad AV. Television viewing and sleep are associated with overweight among urban and semi-urban South Indian children. *Nutr J.* 2007 Sep 20;6:25.
- Landis AM, Parker KP, Dunbar SB. Sleep, hunger, satiety, food cravings, and caloric intake in adolescents. *J Nurs Scholarsh.* 2009;41(2):115-23.
- Lee JA, Park HS. Relation between sleep duration, overweight, and metabolic syndrome in Korean adolescents. *Nutr Metab Cardiovasc Dis.* 2014 Jan;24(1):65-71.
- Lehto R, Ray C, Lahti-Koski M, Roos E. Health behaviors, waist circumference and waist-to-height ratio in children. *Eur J Clin Nutr.* 2011 Jul;65(7):841-8.
- Leprout R, Van Cauter E. Role of sleep and sleep loss in hormonal release and metabolism. *Endocr Dev.* 2010;17:11-21.
- Liu J, Zhang A, Li L. Sleep duration and overweight/obesity in children: review and implications for pediatric nursing. *J Spec Pediatr Nurs.* 2012 Jul;17(3):193-204.
- Liu X, Forbes EE, Ryan ND, Rofey D, Hannon TS, Dahl RE. Rapid eye movement sleep in relation to overweight in children and adolescents. *Arch Gen Psychiatry.* 2008 Aug;65(8):924-32.
- Lumeng JC, Somashekar D, Appugliese D, Kaciroti N, Corwyn RF, Bradley RH. Shorter sleep duration is associated with increased risk for being overweight at ages 9 to 12 years. *Pediatrics.* 2007 Nov;120(5):1020-9.
- Lytte LA, Murray DM, Laska MN, Pasch KE, Anderson SE, Farbaksh K. Examining the longitudinal relationship

- between change in sleep and obesity risk in adolescents. *Health Educ Behav.* 2013 Jun;40(3):362-70.
- Lytle LA, Pasch KE, Farbakhsh K. The relationship between sleep and weight in a sample of adolescents. *Obesity (Silver Spring).* 2011 Feb;19(2):324-31.
- Magee CA, Caputi P, Iverson DC. The longitudinal relationship between sleep duration and body mass index in children: a growth mixture modeling approach. *J Dev Behav Pediatr.* 2013 Apr;34(3):165-73.
- Mahler B, Kamperis K, Schroeder M, Frøkiær J, Djurhuus JC, Rittig S. Sleep deprivation induces excess diuresis and natriuresis in healthy children. *Am J Physiol Renal Physiol.* 2012 Jan 15;302(2):F236-43.
- Martinez SM, Greenspan LC, Butte NF, Gregorich SE, De Groat CL, Deardorff J, Penilla C, Pasch LA, Flores E, Tschann JM. Mother-reported sleep, accelerometer-estimated sleep and weight status in Mexican American children: sleep duration is associated with increased adiposity and risk for overweight/obese status. *J Sleep Res.* 2014 Jun;23(3):326-34.
- Martinez-Gomez D, Eisenmann JC, Gomez-Martinez S, Hill EE, Zapatera B, Veiga OL, Marcos A; AFINOS Study Group. Sleep duration and emerging cardiometabolic risk markers in adolescents. The AFINOS study. *Sleep Med.* 2011 Dec;12(10):997-1002.
- Martinez-Gomez D, Moreno LA, Romeo J, Rey-López P, Castillo R, Cabero MJ, Vicente-Rodriguez G, Gutiérrez A, Veiga OL. Combined influence of lifestyle risk factors on body fat in Spanish adolescents--the Avena study. *Obes Facts.* 2011;4(2):105-11.
- Matthews KA, Dahl RE, Owens JF, Lee L, Hall M. Sleep duration and insulin resistance in healthy black and white adolescents. *Sleep.* 2012 Oct 1;35(10):1353-8.
- Mcneil J, Tremblay MS, Leduc G, Boyer C, Bélanger P, Leblanc AG, Borghese MM, Chaput JP. Objectively-measured sleep and its association with adiposity and physical activity in a sample of Canadian children. *J Sleep Res.* 2015 Apr;24(2):131-9.
- Meininger JC, Gallagher MR, Eissa MA, Nguyen TQ, Chan W. Sleep duration and its association with ambulatory blood pressure in a school-based, diverse sample of adolescents. *Am J Hypertens.* 2014 Jul;27(7):948-55.
- Meng LP, Liu AL, Hu X, Zhang Q, Du SM, Fang HY, Ma J, Xu GF, Li Y, Guo HW, Du L, Ma GS. Report on childhood obesity in China (10): association of sleep duration with obesity. *Biomed Environ Sci.* 2012 Apr;25(2):133-40.
- Michels N, Verbeiren A, Ahrens W, De Henauw S, Sioen I. Children's sleep quality: relation with sleep duration and adiposity. *Public Health.* 2014 May;128(5):488-90.
- Miller AL, Kaciroti N, Lebourgeois MK, Chen YP, Sturza J, Lumeng JC. Sleep timing moderates the concurrent sleep duration-body mass index association in low-income preschool-age children. *Acad Pediatr.* 2014 Mar-Apr;14(2):207-13.
- Mitchell JA, Rodriguez D, Schmitz KH, Audrain-McGovern J. Sleep duration and adolescent obesity. *Pediatrics.* 2013 May;131(5):e1428-34.
- Monasta L, Batty GD, Cattaneo A, Lutje V, Ronfani L, Van Lenthe FJ, Brug J. Early-life determinants of overweight and obesity: a review of systematic reviews. *Obes Rev.* 2010 Oct;11(10):695-708.
- Moraleda-Cibrián M, O'Brien LM. Sleep duration and body mass index in children and adolescents with and without obstructive sleep apnea. *Sleep Breath.* 2014 Sep;18(3):555-61.
- Morley BC, Scully ML, Niven PH, Okely AD, Baur LA, Pratt IS, Wakefield MA; NaSSDA Study Team. What factors are associated with excess body weight in Australian secondary school students? *Med J Aust.* 2012 Feb 20;196(3):189-92.
- Must A, Parisi SM. Sedentary behavior and sleep: paradoxical effects in association with childhood obesity. *Int J Obes (Lond).* 2009 Apr;33 Suppl 1:S82-6.
- Nixon GM, Thompson JM, Han DY, Becroft DM, Clark PM, Robinson E, Waldie KE, Wild CJ, Black PN, Mitchell EA. Short sleep duration in middle childhood: risk factors and consequences. *Sleep.* 2008 Jan;31(1):71-8.
- Nugent R, Althouse A, Yaqub Y, Nugent K, Raj R. Modeling the relation between obesity and sleep parameters in children referred for dietary weight reduction intervention. *South Med J.* 2014 Aug;107(8):473-80.
- O'Dea JA, Dibley MJ, Rankin NM. Low sleep and low socioeconomic status predict high body mass index: a 4-year longitudinal study of Australian schoolchildren. *Pediatr Obes.* 2012 Aug;7(4):295-303.
- Ochiai H, Shirasawa T, Shimada N, Ohtsu T, Nishimura R, Morimoto A, Hoshino H, Tajima N, Kokaze A. Sleep duration and overweight among elementary schoolchildren: a population-based study in Japan. *Acta Med Okayama.* 2012;66(2):93-9.
- Olds T, Blunden S, Dollman J, Maher CA. Day type and the relationship between weight status and sleep duration in children and adolescents. *Aust N Z J Public Health.* 2010 Apr;34(2):165-71.
- Olds TS, Maher CA, Matricciani L. Sleep duration or bedtime? Exploring the relationship between sleep habits and weight status and activity patterns. *Sleep.* 2011 Oct 1;34(10):1299-307.
- Ozturk A, Mazicioglu M, Poyrazoglu S, Cicek B, Gunay O, Kurtoglu S. The relationship between sleep duration and obesity in Turkish children and adolescents. *Acta Paediatr.* 2009 Apr;98(4):699-702.
- Padez C, Mourao I, Moreira P, Rosado V. Long sleep duration and childhood overweight/obesity and body fat. *Am J Hum Biol.* 2009 May-Jun;21(3):371-6.
- Park S. The association between short sleep duration and body mass index among South Korean children and adolescents. *J Sch Nurs.* 2013 Apr;29(2):142-50.
- Park S. Association between short sleep duration and obesity among South Korean adolescents. *West J Nurs Res.* 2011 Mar;33(2):207-23.
- Prats-Puig A, Grau-Cabrera P, Riera-Pérez E, Cortés-Marina R, Fortea E, Soriano-Rodríguez P, de Zegher F, Ibáñez L, Bassols J, López-Bermejo A. Variations in the obesity genes FTO, TMEM18 and NRXN3 influence the vulnerability of children to weight gain induced by short sleep duration. *Int J Obes (Lond).* 2013 Feb;37(2):182-7.
- Reilly JJ, Armstrong J, Dorosty AR, Emmett PM, Ness A, Rogers I, Steer C, Sherriff A; Avon Longitudinal Study of

- Parents and Children Study Team. Early life risk factors for obesity in childhood: cohort study. *BMJ*. 2005 Jun 11;330(7504):1357.
- Reither EN, Krueger PM, Hale L, Reiter EM, Peppard PE. Ethnic variation in the association between sleep and body mass among US adolescents. *Int J Obes (Lond)*. 2014 Jul;38(7):944-9.
- Rey-López JP, de Carvalho HB, de Moraes AC, Ruiz JR, Sjöström M, Marcos A, Polito A, Gottrand F, Manios Y, Kafatos A, Molnar D, Widhalm K, De Henauw S, Moreno LA; HELENA study. Sleep time and cardiovascular risk factors in adolescents: the HELENA (Healthy Lifestyle in Europe by Nutrition in Adolescence) study. *Sleep Med*. 2014 Jan;15(1):104-10.
- Rutters F, Gerver WJ, Nieuwenhuizen AG, Verhoef SP, Westerterp-Plantenga MS. Sleep duration and body-weight development during puberty in a Dutch children cohort. *Int J Obes (Lond)*. 2010 Oct;34(10):1508-14.
- Sallinen BJ, Hassan F, Olszewski A, Maupin A, Hoban TF, Chervin RD, Woolford SJ. Longer weekly sleep duration predicts greater 3-month BMI reduction among obese adolescents attending a clinical multidisciplinary weight management program. *Obes Facts*. 2013;6(3):239-46.
- Sasaki A, Yorifuji T, Iwase T, Komatsu H, Takao S, Doi H. Is There Any Association between TV Viewing and Obesity in Preschool Children in Japan? *Acta Med Okayama*. 2010 Apr;64(2):137-42.
- Seegers V, Petit D, Falissard B, Vitaro F, Tremblay RE, Montplaisir J, Touchette E. Short sleep duration and body mass index: a prospective longitudinal study in preadolescence. *Am J Epidemiol*. 2011 Mar 15;173(6):621-9.
- Seicean A, Redline S, Seicean S, Kirchner HL, Gao Y, Sekine M, Zhu X, Storfer-Isser A. Association between short sleeping hours and overweight in adolescents: results from a US Suburban High School survey. *Sleep Breath*. 2007 Dec;11(4):285-93.
- Sekine M, Yamagami T, Handa K, Saito T, Nanri S, Kawaminami K, Tokui N, Yoshida K, Kagamimori S. A dose-response relationship between short sleeping hours and childhood obesity: results of the Toyama Birth Cohort Study. *Child Care Health Dev*. 2002 Mar;28(2):163-70.
- Shan XY, Xi B, Cheng H, Hou DQ, Wang Y, Mi J. Prevalence and behavioral risk factors of overweight and obesity among children aged 2-18 in Beijing, China. *Int J Pediatr Obes*. 2010 Oct;5(5):383-9.
- Shi Z, Taylor AW, Gill TK, Tuckerman J, Adams R, Martin J. Short sleep duration and obesity among Australian children. *BMC Public Health*. 2010 Oct 15;10:609.
- Shochat T, Cohen-Zion M, Tzischinsky O. Functional consequences of inadequate sleep in adolescents: a systematic review. *Sleep Med Rev*. 2014 Feb;18(1):75-87.
- Silva GE, Goodwin JL, Parthasarathy S, Sherrill DL, Vana KD, Drescher AA, Quan SF. Longitudinal association between short sleep, body weight, and emotional and learning problems in Hispanic and Caucasian children. *Sleep*. 2011 Sep 1;34(9):1197-205.
- Sivertsen B, Pallesen S, Sand L, Hysing M. Sleep and body mass index in adolescence: results from a large population-based study of Norwegian adolescents aged 16 to 19 years. *BMC Pediatr*. 2014 Aug 15;14:204.
- Skidmore PM, Howe AS, Polak MA, Wong JE, Lubransky A, Williams SM, Black KE. Sleep duration and adiposity in older adolescents from Otago, New Zealand: relationships differ between boys and girls and are independent of food choice. *Nutr J*. 2013 Sep 14;12:128.
- Sokolovic N, Kuriyan R, Kurpad AV, Thomas T. Sleep and birthweight predict visceral adiposity in overweight/obese children. *Pediatr Obes*. 2013 Jun;8(3):e41-4.
- Spruyt K, Molfese DL, Gozal D. Sleep duration, sleep regularity, body weight, and metabolic homeostasis in school-aged children. *Pediatrics*. 2011 Feb;127(2):e345-52.
- Steele RM, van Sluijs EM, Cassidy A, Griffin SJ, Ekelund U. Targeting sedentary time or moderate- and vigorous-intensity activity: independent relations with adiposity in a population-based sample of 10-y-old British children. *Am J Clin Nutr*. 2009 Nov;90(5):1185-92.
- Sugimori H, Yoshida K, Izuno T, Miyakawa M, Suka M, Sekine M, Yamagami T, Kagamimori S. Analysis of factors that influence body mass index from ages 3 to 6 years: A study based on the Toyama cohort study. *Pediatr Int*. 2004 Jun;46(3):302-10.
- Sun Y, Sekine M, Kagamimori S. Lifestyle and overweight among Japanese adolescents: the Toyama Birth Cohort Study. *J Epidemiol*. 2009;19(6):303-10.
- Sung V, Beebe DW, Vandyske R, Fenchel MC, Crimmins NA, Kirk S, Hiscock H, Amin R, Wake M. Does sleep duration predict metabolic risk in obese adolescents attending tertiary services? A cross-sectional study. *Sleep*. 2011 Jul 1;34(7):891-8.
- Taheri S. The link between short sleep duration and obesity: we should recommend more sleep to prevent obesity. *Arch Dis Child*. 2006 Nov;91(11):881-4.
- Tan E, Healey D, Gray AR, Galland BC. Sleep hygiene intervention for youth aged 10 to 18 years with problematic sleep: a before-after pilot study. *BMC Pediatr*. 2012 Dec 7;12:189.
- Tatone-Tokuda F, Dubois L, Ramsay T, Girard M, Touchette E, Petit D, Montplaisir JY. Sex differences in the association between sleep duration, diet and body mass index: a birth cohort study. *J Sleep Res*. 2012 Aug;21(4):448-60.
- Taveras EM, Gillman MW, Peña MM, Redline S, Rifas-Shiman SL. Chronic sleep curtailment and adiposity. *Pediatrics*. 2014 Jun;133(6):1013-22.
- Taveras EM, Rifas-Shiman SL, Oken E, Gunderson EP, Gillman MW. Short sleep duration in infancy and risk of childhood overweight. *Arch Pediatr Adolesc Med*. 2008 Apr;162(4):305-11.
- Tian Z, Ye T, Zhang X, Liu E, Wang W, Wang P, Liu G, Yang X, Hu G, Yu Z. Sleep duration and hyperglycemia among obese and nonobese children aged 3 to 6 years. *Arch Pediatr Adolesc Med*. 2010 Jan;164(1):46-52.
- Tikotzky L, DE Marcas G, Har-Toov J, Dollberg S, Bar-Haim Y, Sadeh A. Sleep and physical growth in infants during the first 6 months. *J Sleep Res*. 2010 Mar;19(1 Pt 1):103-10.

- Touchette E, Petit D, Tremblay RE, Boivin M, Falissard B, Genolini C, Montplaisir JY. Associations between sleep duration patterns and overweight/obesity at age 6. *Sleep*. 2008 Nov;31(11):1507-14.
- Vaezghasemi M, Lindkvist M, Ivarsson A, Eurenus E. Overweight and lifestyle among 13-15 year olds: a cross-sectional study in northern Sweden. *Scand J Public Health*. 2012 May;40(3):221-8.
- von Kries R, Toschke AM, Wurmser H, Sauerwald T, Koletzko B. Reduced risk for overweight and obesity in 5- and 6-y-old children by duration of sleep--a cross-sectional study. *Int J Obes Relat Metab Disord*. 2002 May;26(5):710-6.
- Wake M, Price A, Clifford S, Ukoumunne OC, Hiscock H. Does an intervention that improves infant sleep also improve overweight at age 6? Follow-up of a randomised trial. *Arch Dis Child*. 2011 Jun;96(6):526-32.
- Weiss A, Xu F, Storfer-Isser A, Thomas A, Ievers-Landis CE, Redline S. The association of sleep duration with adolescents' fat and carbohydrate consumption. *Sleep*. 2010 Sep;33(9):1201-9.
- Weiss R, Bremer AA, Lustig RH. What is metabolic syndrome, and why are children getting it? *Ann N Y Acad Sci*. 2013 Apr;1281:123-40.
- Wells JC, Hallal PC, Reichert FF, Menezes AM, AraÃjo CL, Victora CG. Sleep patterns and television viewing in relation to obesity and blood pressure: evidence from an adolescent Brazilian birth cohort. *Int J Obes (Lond)*. 2008 Jul;32(7):1042-9.
- Wheaton AG, Perry GS, Chapman DP, Croft JB. Self-reported sleep duration and weight-control strategies among U.S. high school students. *Sleep*. 2013 Aug 1;36(8):1139-45.
- Wing YK, Li SX, Li AM, Zhang J, Kong AP. The effect of weekend and holiday sleep compensation on childhood overweight and obesity. *Pediatrics*. 2009 Nov;124(5):e994-e1000.
- Wong WW, Ortiz CL, Lathan D, Moore LA, Konzelmann KL, Adolph AL, Smith EO, Butte NF. Sleep duration of underserved minority children in a cross-sectional study. *BMC Public Health*. 2013 Jul 12;13:648.
- Zhu Y, Li AM, Au CT, Kong AP, Zhang J, Wong CK, Chan JC, Wing YK. Association between sleep architecture and glucose tolerance in children and adolescents. *J Diabetes*. 2015 Jan;7(1):10-5.